

Projective wellorders and maximal families of orthogonal measures with large continuum

Vera Fischer^{a,1,*}, Sy David Friedman^{a,1}, Asger Törnquist^{a,1}

^a*Kurt Gödel Research Center, University of Vienna, Währinger Strasse 25, A-1090 Vienna, Austria*

Abstract

We study maximal orthogonal families of Borel probability measures on 2^ω (abbreviated m.o. families) and show that there are generic extensions of the constructible universe L in which each of the following holds:

1. There is a Δ_3^1 -definable well order of the reals, there is a Π_2^1 -definable m.o. family, there are no Σ_2^1 -definable m.o. families and $\mathfrak{b} = \mathfrak{c} = \omega_3$ (in fact any reasonable value of \mathfrak{c} will do).
2. There is a Δ_3^1 -definable well order of the reals, there is a Π_2^1 -definable m.o. family, there are no Σ_2^1 -definable m.o. families, $\mathfrak{b} = \omega_1$ and $\mathfrak{c} = \omega_2$.

Keywords: coding, projective wellorders, projective families of orthogonal measures, large continuum

2000 MSC: 03E15, 03E20, 03E35, 03E45

1. Introduction

Let X be a Polish space, and let $P(X)$ denote the Polish space of Borel probability measures on X , in the sense of [9, 17.E]. Recall that if $\mu, \nu \in P(X)$ then μ and ν are said to be *orthogonal*, written $\mu \perp \nu$, if there is a Borel set $B \subseteq X$ such that $\mu(B) = 0$ and $\nu(X \setminus B) = 0$. A set of measures $\mathcal{A} \subseteq P(X)$ is said to be *orthogonal* if whenever $\mu, \nu \in \mathcal{A}$ and $\mu \neq \nu$ then $\mu \perp \nu$. A *maximal orthogonal family*, or *m.o. family*, is an orthogonal family $\mathcal{A} \subseteq P(X)$ which is maximal under inclusion.

The present paper is concerned with the study of *definable* m.o. families. A well-known result to Preiss and Rataj [13] states that there are no analytic m.o. families, and in a recent paper [3] it was shown by Fischer and Törnquist that if all reals are constructible then there is a Π_1^1 m.o. family. The latter paper also raised the question how restrictive the existence of a definable m.o. family

*Corresponding author; Phone: (43-1) 43 1 4277 50523; Fax: (43-1) 43 1 4277 50599.

Email addresses: `vfischer@logic.univie.ac.at` (Vera Fischer), `sdf@logic.univie.ac.at` (Sy David Friedman), `asger@logic.univie.ac.at` (Asger Törnquist)

¹The authors would like to thank the Austrian Science Fund FWF for the generous support through grants no. P 20835-N13 (Fischer, Friedman), and P 19375-N18 (Friedman, Törnquist), as well as a Marie Curie grant from the European Union no. IRG-249167 (Törnquist).

is on the structure of the real line, since it was shown that Π_1^1 m.o. families cannot coexist with Cohen reals.

In the present paper we study Π_2^1 m.o. families in the context of $\mathfrak{c} \geq \aleph_2$, with the additional requirement that there is a Δ_3^1 -definable wellorder of \mathbb{R} . Our main results are:

Theorem 1. It is consistent with $\mathfrak{c} = \mathfrak{b} = \aleph_3$ that there is a Δ_3^1 -definable wellorder of the reals, a Π_2^1 definable maximal orthogonal family of measures and there are no Σ_2^1 -definable maximal sets of orthogonal measures.

There is nothing special about $\mathfrak{c} = \aleph_3$. In fact the same result can be obtained for any reasonable value of \mathfrak{c} .

Theorem 2. It is consistent with $\mathfrak{b} = \aleph_1$, $\mathfrak{c} = \aleph_2$ that there is a Δ_3^1 -definable wellorder of the reals, a Π_2^1 definable maximal orthogonal family of measures and there are no Σ_2^1 -definable maximal sets of orthogonal measures.

Taken together these theorems show that the existence of a Π_2^1 m.o. family does not seem to impose any severe restrictions on the structure of the real line. On the other hand, we show (Proposition 1) that Σ_2^1 m.o. families cannot coexist with neither Cohen nor random reals, which is why in the models produced to prove Theorems 1 and 2 there are no Σ_2^1 m.o. families.

The theorems of this paper belong to a line of results concerning the definability of certain combinatorial objects on the real line and in particular the question of how low in the projective hierarchy such objects exist. In [12] Mathias showed that there is no Σ_1^1 -definable maximal almost disjoint (mad) family in $[\omega]^\omega$. Assuming $V = L$, Miller obtained (see [11]) a Π_1^1 mad family in $[\omega]^\omega$.

The study of the existence of definable combinatorial objects on \mathbb{R} in the presence of a projective wellorder of the reals and $\mathfrak{c} \geq \aleph_2$ was initiated in [1], [4] and [2]. The wellorder of \mathbb{R} in all those models has a Δ_3^1 -definition, which is indeed optimal for models of $\mathfrak{c} \geq \aleph_2$, since by Mansfield's theorem (see [7, Theorem 25.39]) the existence of a Σ_2^1 -definable wellorder of the reals implies that all reals are constructible. The existence of a Π_2^1 -definable ω -mad family in $[\omega]^\omega$ in the presence of $\mathfrak{c} = \mathfrak{b} = \aleph_2$ was established by Friedman and Zdomsky in [4]. In the same paper, referring to earlier results (see [14] and [8]) they outlined the construction of a model in which $\mathfrak{c} = \aleph_2$ and there is a Π_1^1 -definable ω -mad family: start with the constructible universe L , obtain a Π_1^1 -definable ω mad family and proceed with a countable support iteration of length ω_2 of Miller forcing. The techniques were further developed in [2] to establish a model in which there is a Π_2^1 -definable ω -mad family and $\mathfrak{c} = \mathfrak{b} = \aleph_3$. In particular, in the models from [4] and [2], there are no maximal almost disjoint families of size $< \mathfrak{c}$ and so the almost disjointness number has a Π_2^1 -witness.

The present paper combines the encoding techniques of [3] with the techniques of [1, 4, 2] to obtain Theorems 1 and 2. We note that one significant difference from the situation for mad families is that m.o. families always have size \mathfrak{c} (see [3, Proposition 4.1]).

2. Preliminaries

In this section, we briefly recall the coding of probability measures on 2^ω and the encoding technique for measures introduced in [3].

Let X be a Polish space. Recall that measures $\mu, \nu \in P(X)$ then μ is said to be *absolutely continuous* with respect to ν , written $\mu \ll \nu$, if for all Borel subsets of X we have that $\nu(B) = 0$ implies that $\mu(B) = 0$. Two measures $\mu, \nu \in P(2^\omega)$ are called *absolutely equivalent*, written $\mu \approx \nu$, if $\mu \ll \nu$ and $\nu \ll \mu$.

If $s \in 2^{<\omega}$ we let $N_s = \{x \in 2^\omega : s \subseteq x\}$ be the basic neighbourhood determined by s . Following [3], we let

$$p(2^\omega) = \{f : 2^{<\omega} \rightarrow [0, 1] : f(\emptyset) = 1 \wedge (\forall s \in 2^{<\omega}) f(s) = f(s \hat{\ } 0) + f(s \hat{\ } 1)\}.$$

The spaces $p(2^\omega)$ and $P(2^\omega)$ are homeomorphic via the recursively defined isomorphism $f \mapsto \mu_f$ where $\mu_f \in P(2^\omega)$ is the measure uniquely determined by requiring that $\mu_f(N_s) = f(s)$ for all $s \in 2^{<\omega}$. We call the unique real $f \in p(2^\omega)$ such that $\mu = \mu_f$ the *code* for μ . The identification of $P(2^\omega)$ and $p(2^\omega)$ allow us to use the notions of effective descriptive set theory in the space $P(2^\omega)$. For instance, the set $P_c(2^\omega)$ of all non-atomic probability measures on 2^ω is arithmetical because $p_c(2^\omega) = \{f \in p(2^\omega) : \mu_f \text{ is non-atomic}\}$ is easily seen to be arithmetical, as shown in [3].

We will use the method of coding a real $z \in 2^\omega$ into a measure $\mu \in P_c(2^\omega)$ introduced in [3]. For convenience we repeat the construction in minimal detail. Given $\mu \in P_c(2^\omega)$ and $s \in 2^{<\omega}$ we let $t(s, \mu)$ be the lexicographically least $t \in 2^{<\omega}$ such that $s \subseteq t$, $\mu(N_{t \hat{\ } 0}) > 0$ and $\mu(N_{t \hat{\ } 1}) > 0$, if it exists and otherwise we let $t(s, \mu) = \emptyset$. Define recursively $t_n^\mu \in 2^{<\omega}$ by letting $t_0^\mu = \emptyset$ and $t_{n+1}^\mu = t(t_n^\mu \hat{\ } 0, \mu)$. Since μ is non-atomic, we have $\text{lh}(t_{n+1}^\mu) > \text{lh}(t_n^\mu)$. Let $t_\infty^\mu = \bigcup_{n=0}^\infty t_n^\mu$. For $f \in p_c(2^\omega)$ and $n \in \omega \cup \{\infty\}$ we will write t_n^f for $t_n^{\mu_f}$. Clearly the sequence $(t_n^f : n \in \omega)$ is recursive in f .

Define the relation $R \subseteq p_c(2^\omega) \times 2^\omega$ as follows:

$$\begin{aligned} R(f, z) \iff (\forall n \in \omega) (z(n) = 1 \iff (f(t_n^f \hat{\ } 0) = \frac{2}{3}f(t_n^f) \wedge f(t_n^f \hat{\ } 1) = \frac{1}{3}f(t_n^f))) \\ \wedge (z(n) = 0 \iff f(t_n^f \hat{\ } 0) = \frac{1}{3}f(t_n^f) \wedge f(t_n^f \hat{\ } 1) = \frac{2}{3}f(t_n^f)). \end{aligned}$$

Whenever $(f, z) \in R$ we say that f *codes* z . Note that $\text{dom}(R) = \{f \in p_c(2^\omega) : (\exists z) R(f, z)\}$ is Π_1^0 and so the function $r : \text{dom}(R) \rightarrow 2^\omega$, where $r(f) = z$ if and only if $(f, z) \in R$, is also Π_1^0 . If ν is a measure such that $\nu = \mu_f$ for some code f , then let $r(\nu) = r(f)$. The key properties of this construction is contained in the following Lemma (see [3, Coding Lemma]):

Lemma 1. There is a recursive function $G : p_c(2^\omega) \times 2^\omega \rightarrow p_c(2^\omega)$ such that $\mu_{G(f,z)} \approx \mu_f$ and $R(G(f,z), z)$ for all $f \in p_c(2^\omega)$ and $z \in 2^\omega$.

The proofs of Theorems 1 and 2 use the following result, which we now prove.

Proposition 1. Suppose that there either is a Cohen real over L or there is a random real over L . Then there is no Σ_2^1 m.o. family.

We first need a preparatory Lemma. In 2^ω , consider the equivalence E_I defined by

$$xE_I y \iff \sum_{n=0}^{\infty} \frac{|x(n) - y(n)|}{n+1} < \infty.$$

We identify 2^ω with \mathbb{Z}_2^ω and equip it with the Haar measure μ .

Lemma 2. Let $A \subseteq 2^\omega$ be a Borel set such that $\mu(A) > 0$. Then $E_I \leq_B E_I \upharpoonright A$, where $E \upharpoonright A$ is the restriction of E_I to A .

Notation: The constant 0 sequence of length $n \in \omega \cup \{\infty\}$ is denoted 0^n . If $A \subseteq 2^\omega$ and $s \in 2^{<\omega}$ let

$$A_{(s)} = \{x \in 2^\omega : s \hat{\ } x \in A\},$$

the *localization* of A at s .

Proof of Lemma 2. Without loss of generality assume that $A \subseteq 2^\omega$ is closed. We will define $q_n \in \omega$, $s_{n,i}, s_t \in 2^{<\omega}$ recursively for all $n \in \omega$, $i \in \{0, 1\}$ and $t \in 2^{<\omega}$ satisfying

1. $q_0 = 0$ and $q_{n+1} = q_n + \text{lh}(s_{n,0})$.
2. $s_{0,i} = \emptyset$ and $\text{lh}(s_{n,i}) = \text{lh}(s_{n,i-1}) > 0$ when $n > 0$.
3. $s_\emptyset = \emptyset$ and $s_t \hat{\ } i = s_t \hat{\ } s_{\text{lh}(t)+1,i}$ for all $t \in 2^{<\omega}$, $i \in \{0, 1\}$.
4. $\frac{1}{n+1} \leq \sum_{k=0}^{\text{lh}(s_{n+1,0})} \frac{|s_{n+1,0}(k) - s_{n+1,1}(k)|}{q_n + k + 1} \leq \frac{2}{n+1}$.
5. $N_{s_t} \subseteq A$.
6. If $t \in 2^n$ then $\mu(A_{(s_t)}) > 1 - 2^{-n}$.

Suppose this can be done. We claim that the map $2^\omega \rightarrow A : x \mapsto a_x$ defined by

$$a_x = \bigcup_{n \in \omega} s_{x \upharpoonright n}$$

is a Borel (in fact, continuous) reduction of E_I to $E_I \upharpoonright A$. To see this, fix $x, y \in 2^\omega$ and note that by (4) we have that

$$\sum_{n=0}^{\infty} \frac{|x(n) - y(n)|}{n+1} \leq \sum_{n=0}^{\infty} \sum_{k=0}^{\text{lh}(s_{n+1,0})} \frac{|s_{n+1,x(i)}(k) - s_{n+1,y(i)}(k)|}{q_n + k + 1} = \sum_{n=0}^{\infty} \frac{|a_x(n) - a_y(n)|}{n+1} \leq 2 \sum_{n=0}^{\infty} \frac{|x(n) - y(n)|}{n+1}$$

so that $x E_I y$ if and only if $a_x E_I a_y$.

We now show that we can construct a scheme satisfying (1)–(6) above. Suppose $q_k, s_{k,i}$ and s_t have been defined for all $k \leq n$ and $t \in 2^{\leq n}$. It is enough to define $s_{n+1,i}$ satisfying (4)–(6). Define

$$f_{q_n} : 2^\omega \rightarrow [0, \infty] : f_{q_n}(x) = \sum_{k=0}^{\infty} \frac{x(k)}{q_n + k + 1}.$$

It is clear that $f_{q_n}(N_{0^k})$ is dense in $[0, \infty]$ for all $k \in \omega$. Let

$$A' = \{x \in A : \lim_{k \rightarrow \infty} \mu(A_{(x \upharpoonright k)}) \rightarrow 1\},$$

i.e, the set of points in A of density 1. By the Lebesgue density theorem [9, 17.9] we have $\mu(A \setminus A') = 0$. Let $A'' = \bigcap_{t \in 2^n} A'_{(s_t)}$ and note that by (6) we have $\mu(A'') > 0$. Thus the set of differences $A'' - A''$ contains a neighborhood of 0^∞ by [9, 17.13]. It follows that there are $x_0, x_1 \in A''$ such that

$$\frac{1}{n+2} \leq \sum_{k=0}^{\infty} \frac{|x_0(k) - x_1(k)|}{q_n + k + 1} \leq \frac{2}{n+2}.$$

Since all points in $A'_{(s_t)}$ have density 1 in $A'_{(s_t)}$ there is some $k_0 \in \omega$ such that

$$\mu(A'_{(s_t \upharpoonright x_i \upharpoonright k_0)}) > 1 - 2^{-n-1}$$

for all $t \in 2^n$. Defining $s_{n+1,i} = x_i \upharpoonright k_0$, it is then clear that (4)–(6) holds. \square

Proof of Proposition 1. We proceed exactly as in [3, Proposition 4.2]. Suppose $A \subseteq P(2^\omega)$ is a Σ_2^1 m.o. family. Recall from [10] and [3, p. 1406] that there is a Borel function $2^\omega \rightarrow P(2^\omega) : x \mapsto \mu^x$ such that

$$xE_I y \implies \mu^x \approx \mu^y$$

and

$$x \not E_I y \implies \mu^x \perp \mu^y.$$

Define as in [3, Proposition 4.2] a relation $Q \subseteq 2^\omega \times P(2^\omega)^\omega$ by

$$Q(x, (\nu_n)) \iff (\forall n)(\nu_n \in A \wedge \nu_n \not\perp \mu^x) \wedge (\forall \mu)(\mu \not\perp \mu^x \implies (\exists n)\nu_n \not\perp \mu)$$

and note that this is Σ_2^1 when A is. Note that $Q(x, (\nu_n))$ precisely when (ν_n) enumerates the measures in A not orthogonal to μ^x (this set is always countable, see [10, Theorem 3.1].) Since A is maximal, each section Q_x is non-empty, and so we can uniformize Q with a (total) function $f : 2^\omega \rightarrow p(2^\omega)^\omega$ having a Δ_2^1 graph. Note that assignment

$$x \mapsto A(x) = \{f(x)_n : n \in \mathcal{N}\}$$

is invariant on the E_I classes.

If there is a Cohen real over L it follows from [6] that f is Baire measurable. Since E_I is a turbulent equivalence relation (in the sense of Hjorth, see e.g. [10]) the map $x \mapsto A(x)$ must be constant on a comeagre set. But this contradicts that all E_I classes are meagre.

If on the other hand there is a random real over L , then f is Lebesgue measurable by [6]. Let $F \subseteq 2^\omega$ be a closed set with positive measure on which f is continuous, and let $g : 2^\omega \rightarrow F$ be a Borel reduction of E_I to $E_I \upharpoonright F$. Note that $x \mapsto A(g(x))$ is then an E_I -invariant Borel assignment of countable subsets of $p(2^\omega)$, and so since E_I is turbulent the function $f \circ g$ must be constant on a comeagre set. This again contradicts that all E_I classes are meagre. \square

3. Δ_3^1 w.o. of the reals, Π_2^1 m.o. family, no Σ_2^1 m.o. families with $\mathfrak{b} = \mathfrak{c} = \aleph_3$

We proceed with the proof of Theorem 1. We will use a modification of the model constructed in [2]. The preliminary stage $\mathbb{P}_0 = \mathbb{P}^0 * \mathbb{P}^1 * \mathbb{P}^2$ of the iteration will coincide almost identically

with the preliminary stage \mathbb{P}_0 of [2] (see Step 0 through Step 2). For convenience of the reader we outline its construction. We work over the constructible universe L .

Recall that a transitive ZF^- model is *suitable* if $\omega_3^{\mathcal{M}}$ exists and $\omega_3^{\mathcal{M}} = \omega_3^{L^{\mathcal{M}}}$. If \mathcal{M} is suitable then also $\omega_1^{\mathcal{M}} = \omega_1^{L^{\mathcal{M}}}$ and $\omega_2^{\mathcal{M}} = \omega_2^{L^{\mathcal{M}}}$.

Fix a $\diamond_{\omega_2}(\text{cof}(\omega_1))$ sequence $\langle G_\xi : \xi \in \omega_2 \cap \text{cof}(\omega_1) \rangle$ which is Σ_1 -definable over L_{ω_2} . For $\alpha < \omega_3$, let W_α be the L -least subset of ω_2 coding α and let $S_\alpha = \{\xi \in \omega_2 \cap \text{cof}(\omega_1) : G_\xi = W_\alpha \cap \xi \neq \emptyset\}$. Then $\vec{S} = \langle S_\alpha : 1 < \alpha < \omega_3 \rangle$ is a sequence of stationary subsets of $\omega_2 \cap \text{cof}(\omega_1)$, which are mutually almost disjoint.

For every α such that $\omega \leq \alpha < \omega_3$ shoot a club C_α disjoint from S_α via the poset \mathbb{P}_α^0 , consisting of all closed subsets of ω_2 which are disjoint from S_α with extension relation end-extension and let $\mathbb{P}^0 = \prod_{\alpha < \omega_3} \mathbb{P}_\alpha^0$ be the direct product of the \mathbb{P}_α^0 's with supports of size ω_1 , where for $\alpha \in \omega$, \mathbb{P}_α^0 is the trivial poset. Then \mathbb{P}^0 is countably closed, ω_2 -distributive and ω_3 -c.c.

For every α such that $\omega \leq \alpha < \omega_3$ let $D_\alpha \subseteq \omega_3$ be a set coding the triple $\langle C_\alpha, W_\alpha, W_\gamma \rangle$ where γ is the largest limit ordinal $\leq \alpha$. Let

$$E_\alpha = \{\mathcal{M} \cap \omega_2 : \mathcal{M} \prec L_{\alpha+\omega_2+1}[D_\alpha], \omega_1 \cup \{D_\alpha\} \subseteq \mathcal{M}\}.$$

Then E_α is a club on ω_2 . Choose $Z_\alpha \subseteq \omega_2$ such that $\text{Even}(Z_\alpha) = D_\alpha$, where $\text{Even}(Z_\alpha) = \{\beta : 2 \cdot \beta \in Z_\alpha\}$, and if $\beta < \omega_2$ is the $\omega_2^{\mathcal{M}}$ for some suitable model \mathcal{M} such that $Z_\alpha \cap \beta \in \mathcal{M}$, then $\beta \in E_\alpha$. Then we have:

(*) $_\alpha$: If $\beta < \omega_2$, \mathcal{M} is a suitable model such that $\omega_1 \subset \mathcal{M}$, $\omega_2^{\mathcal{M}} = \beta$, and $Z_\alpha \cap \beta \in \mathcal{M}$, then $\mathcal{M} \models \psi(\omega_2, Z_\alpha \cap \beta)$, where $\psi(\omega_2, X)$ is the formula “ $\text{Even}(X)$ codes a triple $\langle \bar{C}, \bar{W}, \bar{W} \rangle$, where \bar{W} and \bar{W} are the L -least codes of ordinals $\bar{\alpha}, \bar{\alpha} < \omega_3$ such that $\bar{\alpha}$ is the largest limit ordinal not exceeding $\bar{\alpha}$, and \bar{C} is a club in ω_2 disjoint from $S_{\bar{\alpha}}$ ”.

Similarly to \vec{S} define a sequence $\vec{A} = \langle A_\xi : \xi < \omega_2 \rangle$ of stationary subsets of ω_1 using the “standard” \diamond -sequence. Code Z_α by a subset X_α of ω_1 with the poset \mathbb{P}_α^1 consisting of all pairs $\langle s_0, s_1 \rangle \in [\omega_1]^{<\omega_1} \times [Z_\alpha]^{<\omega_1}$ where $\langle t_0, t_1 \rangle \leq \langle s_0, s_1 \rangle$ iff s_0 is an initial segment of t_0 , $s_1 \subseteq t_1$ and $t_0 \setminus s_0 \cap A_\xi = \emptyset$ for all $\xi \in s_1$. Then X_α satisfies the following condition:

(**) $_\alpha$: If $\omega_1 < \beta \leq \omega_2$ and \mathcal{M} is a suitable model such that $\omega_2^{\mathcal{M}} = \beta$ and $\{X_\alpha\} \cup \omega_1 \subset \mathcal{M}$, then $\mathcal{M} \models \phi(\omega_1, \omega_2, X_\alpha)$, where $\phi(\omega_1, \omega_2, X)$ is the formula: “Using the sequence \vec{A} , X almost disjointly codes a subset \bar{Z} of ω_2 , such that $\text{Even}(\bar{Z})$ codes a triple $\langle \bar{C}, \bar{W}, \bar{W} \rangle$, where \bar{W} and \bar{W} are the L -least codes of ordinals $\bar{\alpha}, \bar{\alpha} < \omega_3$ such that $\bar{\alpha}$ is the largest limit ordinal not exceeding $\bar{\alpha}$, and \bar{C} is a club in ω_2 disjoint from $S_{\bar{\alpha}}$ ”.

Let $\mathbb{P}^1 = \prod_{\alpha < \omega_3} \mathbb{P}_\alpha^1$, where \mathbb{P}_α^1 is the trivial poset for all $\alpha \in \omega$, with countable support. Then \mathbb{P}^1 is countably closed and has the ω_2 -c.c.

Finally we force a localization of the X_α 's. Fix ϕ as in (**) $_\alpha$ and let $\mathcal{L}(X, X')$ be the poset defined in [2, Definition 1], where $X, X' \subset \omega_1$ are such that $\phi(\omega_1, \omega_2, X)$ and $\phi(\omega_1, \omega_2, X')$ hold in any suitable model \mathcal{M} with $\omega_1^{\mathcal{M}} = \omega_1^L$ containing X and X' , respectively. That is $\mathcal{L}(X, X')$ consists of all functions $r : |r| \rightarrow 2$, where the domain $|r|$ of r is a countable limit ordinal such that:

1. if $\gamma < |r|$ then $\gamma \in X$ iff $r(3\gamma) = 1$
2. if $\gamma < |r|$ then $\gamma \in X'$ iff $r(3\gamma + 1) = 1$
3. if $\gamma \leq |r|$, \mathcal{M} is a countable suitable model containing $r \upharpoonright \gamma$ as an element and $\gamma = \omega_1^{\mathcal{M}}$, then $\mathcal{M} \models \phi(\omega_1, \omega_2, X \cap \gamma) \wedge \phi(\omega_1, \omega_2, X' \cap \gamma)$.

The extension relation is end-extension. Then let $\mathbb{P}_{\alpha+m}^2 = \mathcal{L}(X_{\alpha+m}, X_\alpha)$ for every $\alpha \in \text{Lim}(\omega_3) \setminus \{0\}$ and $m \in \omega$. Let $\mathbb{P}_{\alpha+m}^2$ be the trivial poset for $\alpha = 0$, $m \in \omega$ and let

$$\mathbb{P}^2 = \prod_{\alpha \in \text{Lim}(\omega_3)} \prod_{m \in \omega} \mathbb{P}_{\alpha+m}^2$$

with countable supports. Note that the poset $\mathbb{P}_{\alpha+m}^2$, where $\alpha > 0$, produces a generic function in ${}^{\omega_1}2$ (of $L^{\mathbb{P}^0 * \mathbb{P}^1}$), which is the characteristic function of a subset $Y_{\alpha+m}$ of ω_1 with the following property:

$(***)_\alpha$: For every $\beta < \omega_1$ and any suitable \mathcal{M} such that $\omega_1^{\mathcal{M}} = \beta$ and $Y_{\alpha+m} \cap \beta$ belongs to \mathcal{M} , we have $\mathcal{M} \models \phi(\omega_1, \omega_2, X_{\alpha+m} \cap \beta) \wedge \phi(\omega_1, \omega_2, X_\alpha \cap \beta)$.

Claim. $\mathbb{P}_0 := \mathbb{P}^0 * \mathbb{P}^1 * \mathbb{P}^2$ is ω -distributive.

Proof. [2, Lemma 1]. □

Let $\vec{B} = \langle B_{\zeta, m} : \zeta < \omega_1, m \in \omega \rangle$ be a nicely definable sequence of almost disjoint subsets of ω . We will define a finite support iteration $\langle \mathbb{P}_\alpha, \dot{Q}_\beta : \alpha \leq \omega_3, \beta < \omega_3 \rangle$ such that $\mathbb{P}_0 = \mathbb{P}^0 * \mathbb{P}^1 * \mathbb{P}^2$, for every $\alpha < \omega_3$, \dot{Q}_α is a \mathbb{P}_α -name for a σ -centered poset, in $L^{\mathbb{P}^{\omega_3}}$ there is a Δ_3^1 -definable wellorder of the reals, a Π_2^1 -definable maximal family of orthogonal measures and there are no Σ_2^1 -definable maximal families of orthogonal measures. Along the iteration for every $\alpha < \omega_3$, we will define in $V^{\mathbb{P}^\alpha}$ a set O_α of orthogonal measures and for $\alpha \in \text{Lim}(\alpha)$ a subset A_α of $[\alpha, \alpha + \omega)$. Every \mathbb{Q}_α will add a generic real, whose \mathbb{P}_α -name will be denoted \dot{u}_α and similarly to the proof of [2, Lemma 2] one can prove that $L[G_\alpha] \cap {}^\omega\omega = L[\langle \dot{u}_\xi^{G_\alpha} : \xi < \alpha \rangle] \cap {}^\omega\omega$ for every \mathbb{P}_α -generic filter G_α . This gives a canonical wellorder of the reals in $L[G_\alpha]$ which depends only on the sequence $\langle \dot{u}_\xi : \xi < \alpha \rangle$, whose \mathbb{P}_α -name will be denoted by $\dot{<}_\alpha$. We can additionally arrange that for $\alpha < \beta$, $\dot{<}_\alpha$ is an initial segment of $\dot{<}_\beta$, where $\dot{<}_\alpha = \dot{<}_\alpha^{G_\alpha}$ and $\dot{<}_\beta = \dot{<}_\beta^{G_\beta}$. Then if G is a \mathbb{P}_{ω_3} -generic filter over L , then $\dot{<}^G = \bigcup \{ \dot{<}_\alpha^G : \alpha < \omega_3 \}$ will be the desired wellorder of the reals and $O = \bigcup_{\alpha < \omega_3} O_\alpha$ will be the Π_2^1 -definable maximal family of orthogonal measures.

We proceed with the recursive definition of \mathbb{P}_{ω_3} . For every $\nu \in [\omega_2, \omega_3)$ let $i_\nu : \nu \cup \{ \langle \xi, \eta \rangle : \xi < \eta < \nu \} \rightarrow \text{Lim}(\omega_3)$ be a fixed bijection. If G_α is a \mathbb{P}_α -generic filter over L , $\dot{<}_\alpha = \dot{<}_\alpha^{G_\alpha}$ and x, y are reals in $L[G_\alpha]$ such that $x <_\alpha y$, let $x * y = \{ 2n : n \in x \} \cup \{ 2n + 1 : n \in y \}$ and $\Delta(x * y) = \{ 2n + 2 : n \in x * y \} \cup \{ 2n + 1 : n \notin x * y \}$. Suppose \mathbb{P}_α has been defined and fix a \mathbb{P}_α -generic filter G_α .

If $\alpha = \omega_2 \cdot \alpha' + \xi$, where $\alpha' > 0$, $\xi \in \text{Lim}(\omega_2)$, let $\nu = o.t.(\dot{<}_{\omega_2 \cdot \alpha'}^G)$ and let $i = i_\nu$.

Case 1. If $i^{-1}(\xi) = \langle \xi_0, \xi_1 \rangle$ for some $\xi_0 < \xi_1 < \nu$, let x_{ξ_0} and x_{ξ_1} be the ξ_0 -th and ξ_1 -th reals in $L[G_{\omega_2 \cdot \alpha'}]$ according to the wellorder $\dot{<}_{\omega_2 \cdot \alpha'}^{G_\alpha}$. In $L^{\mathbb{P}_\alpha}$ let

$$\mathbb{Q}_\alpha = \{ \langle s_0, s_1 \rangle : s_0 \in [\omega]^{<\omega}, s_1 \in [\bigcup_{m \in \Delta(x_{\xi_0} * x_{\xi_1})} Y_{\alpha+m} \times \{m\}]^{<\omega} \},$$

where $\langle t_0, t_1 \rangle \leq \langle s_0, s_1 \rangle$ if and only if $s_1 \subseteq t_1$, s_0 is an initial segment of t_0 and $(t_0 \setminus s_0) \cap B_{\zeta, m} = \emptyset$ for all $\langle \zeta, m \rangle \in s_1$. Let u_α be the generic real added by \mathbb{Q}_α , $A_\alpha = \alpha + \omega \setminus \Delta(x_{\xi_0} * x_{\xi_1})$ and $O_\alpha = \emptyset$.

Case 2. Suppose $i^{-1}(\xi) = \zeta \in \nu$. If the ζ -th real according to the wellorder $\dot{<}_{\omega_2 \cdot \alpha'}^{G_\alpha}$ is not the code of a measure orthogonal to $O'_\alpha = \bigcup_{\gamma < \alpha} O_\gamma$, let \mathbb{Q}_α be the trivial poset, $A_\alpha = \emptyset$, $O_\alpha = \emptyset$. Otherwise, i.e. in case x_ζ is a code for a measure orthogonal to O'_α , let

$$\mathbb{Q}_\alpha = \{ \langle s_0, s_1 \rangle : s_0 \in [\omega]^{<\omega}, s_1 \in [\bigcup_{m \in \Delta(x_\zeta)} Y_{\alpha+m} \times \{m\}]^{<\omega} \},$$

where $\langle t_0, t_1 \rangle \leq \langle s_0, s_1 \rangle$ if and only if $s_1 \subseteq t_1$, s_0 is an initial segment of t_0 and $(t_0 \setminus s_0) \cap B_{\zeta, m} = \emptyset$ for all $\langle \zeta, m \rangle \in s_1$. Let u_α be the generic real added by \mathbb{Q}_α . In $L^{\mathbb{P}_{\alpha+1}} = L^{\mathbb{P}_\alpha * \mathbb{Q}_\alpha}$ let $g_\alpha = G(x_\zeta, u_\alpha)$ be the code of a measure equivalent to μ_{x_ζ} which codes u_α (see [3, Lemma 3.5]) and let $O_\alpha = \{ \mu_{g_\alpha} \}$. Let $A_\alpha = \alpha + \omega \setminus \Delta(u_\alpha)$.

If α is not of the above form, i.e. α is a successor or $\alpha \in \omega_2$, let \mathbb{Q}_α be the following poset for adding a dominating real:

$$\mathbb{Q}_\alpha = \{ \langle s_0, s_1 \rangle : s_0 \in \omega^{<\omega}, s_1 \in [\text{o.t.}(\dot{<}_\alpha^{G_\alpha})]^{<\omega} \},$$

where $\langle t_0, t_1 \rangle \leq \langle s_0, s_1 \rangle$ if and only if s_0 is an initial segment of t_0 , $s_1 \subseteq t_1$, and $t_0(n) > x_\xi(n)$ for all $n \in \text{dom}(t_0) \setminus \text{dom}(s_0)$ and $\xi \in s_1$, where x_ξ is the ξ -th real in $L[G_\alpha] \cap \omega^\omega$ according to the wellorder $\dot{<}_\alpha^{G_\alpha}$. Let $A_\alpha = \emptyset$, $O_\alpha = \emptyset$.

With this the definition of \mathbb{P}_{ω_3} is complete. Let $O = \bigcup_{\alpha < \omega_3} O_\alpha$. In $L^{\mathbb{P}_{\omega_3}}$ we have: ν is a measure in the set O if and only if for every countable suitable model \mathcal{M} such that $\nu \in \mathcal{M}$, there is $\bar{\alpha} < \omega_3^{\mathcal{M}}$ such that $S_{\bar{\alpha}+m}$ is nonstationary in $(L[r(\nu)])^{\mathcal{M}}$ for every $m \in \Delta(r(\nu))$. Therefore O has indeed a Π_2^1 definition. Furthermore O is maximal in $P_c(2^\omega)$. Indeed, suppose in $L^{\mathbb{P}_{\omega_3}}$ there is a code x for a measure orthogonal to every measure in the family O . Choose α minimal such that $\alpha = \omega_2 \cdot \alpha' + \xi$ for some $\alpha' > 0$ and $\xi \in \text{Lim}(\omega_2)$ and $x \in L[G_{\omega_2 \cdot \alpha'}]$. Let $\nu = \text{o.t.}(\dot{<}_{\omega_2 \cdot \alpha'}^{G_\alpha})$ and let $i = i_\nu$. Then $x = x_\zeta$ is the ζ -th real according to the wellorder $\dot{<}_{\omega_2 \cdot \alpha'}^{G_\alpha}$, where $\zeta \in \nu$ and so for some $\xi \in \text{Lim}(\omega_2)$, $i^{-1}(\xi) = \zeta$. But then $x_\zeta = x$ is the code of a measure orthogonal to O_α and so by construction $O_{\alpha+1}$ contains a measure equivalent to μ_x , which is a contradiction. To obtain a Π_2^1 -definable m.o. family in $L^{\mathbb{P}_{\omega_3}}$ consider the union of O with the set of all point measures. Just as in [2] one can show that $<$ is indeed a Δ_3^1 -definable wellorder of the reals.

Since \mathbb{P}_{ω_3} is a finite support iteration, along the iteration cofinally often we have added Cohen reals. Thus for every real a in $L^{\mathbb{P}_{\omega_3}}$ there is a Cohen real over $L[a]$ and so by Proposition 1 in $L^{\mathbb{P}_{\omega_3}}$ there are no Σ_2^1 m.o. families. Also note that since cofinally often we have added dominating reals, $L^{\mathbb{P}_{\omega_3}} \models \mathfrak{b} = \omega_3$.

4. Δ_3^1 w.o. of the reals, a Π_2^1 m.o. family, no Σ_2^1 m.o. families with $\mathfrak{c} = \aleph_2$

In this section we establish the proof of Theorem 2. The model is obtained as a slight modification of the iteration construction developed in [1]. We restate the definitions of the posets used in this construction. For a more detailed account of their properties see [1]. We work over the constructible universe L .

If $S \subseteq \omega_1$ is a stationary, co-stationary set, then by $Q(S)$ denote the poset of all countable closed subsets of $\omega_1 \setminus S$ with extension relation end-extension. Recall that $Q(S)$ is $\omega_1 \setminus S$ -proper, ω -distributive and adds a club disjoint from S (see [1], [5]). For the proof of Theorem 2 we use the form of localization defined in [1, Definition 1]. That is, if $X \subseteq \omega_1$ and $\phi(\omega_1, X)$ is a Σ_1 -sentence with parameters ω_1, X which is true in all suitable models containing ω_1 and X as elements, then $\mathcal{L}(\phi)$ be the poset of all functions $r : |r| \rightarrow 2$, where the domain $|r|$ of r is a countable limit ordinal, such that

1. if $\gamma < |r|$ then $\gamma \in X$ iff $r(2\gamma) = 1$
2. if $\gamma \leq |r|$, \mathcal{M} is a countable, suitable model containing $r \upharpoonright \gamma$ as an element and $\gamma = \omega_1^{\mathcal{M}}$, then $\phi(\gamma, X \cap \gamma)$ holds in \mathcal{M} .

The extension relation is end-extension. Recall that $\mathcal{L}(\phi)$ has a countably closed dense subset (see [1, Remark 2]) and that if G is $\mathcal{L}(\phi)$ -generic and \mathcal{M} is a countable suitable model containing $(\bigcup G) \upharpoonright \gamma$ as an element, where $\gamma = \omega_1^{\mathcal{M}}$, then $\mathcal{M} \models \phi(\gamma, X \cap \gamma)$ (see [1, Lemma 2]).

We will use also the coding with perfect trees defined in [1, Definition 2]. Let $Y \subseteq \omega_1$ be generic over L such that in $L[Y]$ cofinalities have not been changed and let $\bar{\mu} = \{\mu_i\}_{i \in \omega_1}$ be a sequence of L -countable ordinals such that μ_i is the least $\mu > \sup_{j < i} \mu_j$, $L_{\mu_i}[Y \cap i] \models ZF^-$ and $L_{\mu_i} \models \omega$ is the largest cardinal. Say that a real R codes Y below i if for all $j < i$, $j \in Y$ if and only if $L_{\mu_j}[Y \cap j, R] \models ZF^-$. For $T \subseteq 2^{<\omega}$ a perfect tree, let $|T|$ be the least i such that $T \in L_{\mu_i}[Y \cap i]$. Then $\mathcal{C}(Y)$ is the poset of all perfect trees T such that R codes Y below $|T|$, whenever R is a branch through T , where for T_0, T_1 conditions in $\mathcal{C}(Y)$, $T_0 \leq T_1$ if and only if T_0 is a subtree of T_1 . Recall also that $\mathcal{C}(Y)$ is proper and ${}^\omega\omega$ -bounding (see [1, Lemmas 7,8]).

Fix a bookkeeping function $F : \omega_2 \rightarrow L_{\omega_2}$ and a sequence $\vec{S} = (S_\beta : \beta < \omega_2)$ of almost disjoint stationary subsets of ω_1 , defined as in [1, Lemma 14]. Thus F and \vec{S} are Σ_1 -definable over L_{ω_2} with parameter ω_1 , $F^{-1}(a)$ is unbounded in ω_2 for every $a \in L_{\omega_2}$ and whenever \mathcal{M}, \mathcal{N} are suitable models such that $\omega_1^{\mathcal{M}} = \omega_1^{\mathcal{N}}$ then $F^{\mathcal{M}}, \vec{S}^{\mathcal{M}}$ agree with $F^{\mathcal{N}}, \vec{S}^{\mathcal{N}}$ on $\omega_2^{\mathcal{M}} \cap \omega_2^{\mathcal{N}}$. Also if \mathcal{M} is suitable and $\omega_1^{\mathcal{M}} = \omega_1$ then $F^{\mathcal{M}}, \vec{S}^{\mathcal{M}}$ equal the restrictions of F, \vec{S} to the ω_2 of \mathcal{M} . Fix also a stationary subset S of ω_1 which is almost disjoint from every element of \vec{S} .

Recursively we will define a countable support iteration $\langle \mathbb{P}_\alpha, \dot{Q}_\beta : \alpha \leq \omega_2, \beta < \omega_2 \rangle$ and a sequence $\langle O_\alpha : \alpha \in \omega_2 \rangle$, such that in $L^{\mathbb{P}_{\omega_2}}$ there is a Δ_3^1 -definable wellorder of the reals and $O = \bigcup_{\alpha < \omega_2} O_\alpha$ is a maximal family of orthogonal measures. Define the wellorder $<_\alpha$ in $L[G_\alpha]$ where G_α is \mathbb{P}_α -generic just as in [1]. We can assume that all names for reals are nice and that for $\alpha < \beta < \omega_2$, all \mathbb{P}_α -names for reals precede in the canonical wellorder $<_L$ of L all \mathbb{P}_β -names for reals, which are not \mathbb{P}_α -names. For each $\alpha < \omega_2$, define a wellorder $<_\alpha$ on the reals of $L[G_\alpha]$, where G_α is a \mathbb{P}_α -generic as follows. If x is a real in $L[G_\alpha]$ let σ_x^α be the $<_L$ -least \mathbb{P}_γ -name for x ,

where $\gamma \leq \alpha$ is least so that x has a \mathbb{P}_γ -name. For x, y reals in $L[G_\alpha]$ define $x <_\alpha y$ if and only if $\sigma_x^\alpha <_L \sigma_y^\alpha$. Note that whenever $\alpha < \beta$, then $<_\alpha$ is an initial segment of $<_\beta$.

We proceed with the definition of the poset. Let \mathbb{P}_0 be the trivial poset. Suppose \mathbb{P}_α and $\langle O_\gamma : \gamma < \alpha \rangle$ have been defined. Let $\dot{\mathbb{Q}}_\alpha = \dot{\mathbb{Q}}_\alpha^0 * \dot{\mathbb{Q}}_\alpha^1$ be a \mathbb{P}_α -name for a poset where $\dot{\mathbb{Q}}_\alpha^0$ is a \mathbb{P}_α -name for the random real forcing and $\dot{\mathbb{Q}}_\alpha^1$ is defined as follows:

Case 1. If $F(\alpha) = \{\sigma_x^\alpha, \sigma_y^\alpha\}$ for some pair of reals x, y in $L[G_\alpha]$, then define \mathbb{Q}_α as in [1]. That is \mathbb{Q}_α is a three stage iteration $\mathbb{K}_\alpha^0 * \mathbb{K}_\alpha^1 * \mathbb{K}_\alpha^2$ where:

(1) In $V^{\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha^0}$, \mathbb{K}_α^0 is the direct limit $\langle \mathbb{P}_{\alpha, n}^0, \dot{\mathbb{K}}_{\alpha, n}^0 : n \in \omega \rangle$, where $\dot{\mathbb{K}}_{\alpha, n}^0$ is a $\mathbb{P}_{\alpha, n}^0$ -name for $Q(S_{\alpha+2n})$ for $n \in x_\alpha * y_\alpha$, and $\dot{\mathbb{K}}_{\alpha, n}^0$ is a $\mathbb{P}_{\alpha, n}^0$ -name for $Q(S_{\alpha+2n+1})$ for $n \notin x_\alpha * y_\alpha$.

(2) Let G_α^0 be a $\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha^0$ -generic filter and let H_α be a \mathbb{K}_α^0 -generic over $L[G_\alpha^0]$. In $L[G_\alpha^0 * H_\alpha]$ let X_α be a subset of ω_1 coding α , coding the pair (x_α, y_α) , coding a level of L in which α has size at most ω_1 and coding the generic $G_\alpha^0 * H_\alpha$, which we can regard as a subset of an element of L_{ω_2} . Let $\mathbb{K}_\alpha^1 = \mathcal{L}(\phi_\alpha)$ where $\phi_\alpha = \phi_\alpha(\omega_1, X)$ is the Σ_1 -sentence which holds if and only if X codes an ordinal $\bar{\alpha} < \omega_2$ and a pair (x, y) such that $S_{\bar{\alpha}+2n}$ is nonstationary for $n \in x * y$ and $S_{\bar{\alpha}+2n+1}$ is nonstationary for $n \notin x * y$. Let \dot{X}_α be a $\mathbb{P}_\alpha^0 * \dot{\mathbb{Q}}_\alpha^0 * \dot{\mathbb{K}}_\alpha^0$ -name for X_α and let $\dot{\mathbb{K}}_\alpha^1$ be a $\mathbb{P}_\alpha^0 * \dot{\mathbb{Q}}_\alpha^0 * \dot{\mathbb{K}}_\alpha^0$ -name for \mathbb{K}_α^1 .

(3) Let Y_α be \mathbb{K}_α^1 -generic over $L[G_\alpha^0 * H_\alpha]$. Note that the even part of Y_α -codes X_α and so codes the generic $G_\alpha^0 * H_\alpha$. Then in $L[Y_\alpha] = L[G_\alpha^0 * H_\alpha * Y_\alpha]$, let $\mathbb{K}_\alpha^2 = \mathcal{C}(Y_\alpha)$. Finally, let $\dot{\mathbb{K}}_\alpha^2$ be a $\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha^0 * \dot{\mathbb{K}}_\alpha^0 * \dot{\mathbb{K}}_\alpha^1$ -name for \mathbb{K}_α^2 .

Case 2. If $F(\alpha) = \{\sigma_x^\alpha\}$ where x is a code for a measure orthogonal to $\bigcup_{\gamma < \alpha} O_\gamma$, then let $\dot{\mathbb{Q}}_\alpha^1$ be a $\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha^0$ -name for $\mathbb{K}_\alpha^0 * \mathbb{K}_\alpha^1 * \mathbb{K}_\alpha^2$ where in $L^{\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha^0}$, \mathbb{K}_α^0 is the direct limit $\langle \mathbb{P}_{\alpha, n}^0, \dot{\mathbb{Q}}_{\alpha, n}^0 : n \in \omega \rangle$ where $\dot{\mathbb{Q}}_{\alpha, n}^0$ is a $\mathbb{P}_{\alpha, n}^0$ -name for $Q(S_{\alpha+2n})$ for every $n \in x$ and a $\mathbb{P}_{\alpha, n}^0$ -name for $Q(S_{\alpha+2n+1})$ for every $n \notin x$. Define \mathbb{K}_α^1 and \mathbb{K}_α^2 just as in *Case 1*. In $L^{\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha^0}$ let $g = G(x, R_\alpha)$ be a code for a measure which is equivalent to μ_x and codes the real R_α . Let $O_\alpha = \{\mu_g\}$.

In any other case, let \mathbb{Q}_α be a \mathbb{P}_α -name for the trivial poset, $O_\alpha = \emptyset$. With this the definition of \mathbb{P}_{ω_2} and the family $O = \bigcup_{\gamma < \omega_2} O_\alpha$ is complete.

Claim. $O = \bigcup_{\gamma < \omega_2} O_\gamma$ is a maximal family of orthogonal measures in $P_c(2^\omega)$.

Proof. It is clear that O is a family of orthogonal measures. It remains to verify its maximality. Suppose the contrary and let f be a code for a measure in $L[G]$ where G is \mathbb{P}_{ω_3} -generic over L , which is orthogonal to all measures in O . Fix α minimal such that f is in $L[G_\alpha]$ and let σ be the $<_L$ -least name for f . Since $F^{-1}(\sigma)$ is unbounded, there is $\beta \geq \alpha$ such that $F(\beta) = \{\sigma\}$. Therefore \mathbb{Q}_β is nontrivial and $O_\beta = \{\mu_g\}$ for some measure μ_g which is equivalent to μ_f , which is a contradiction. \square

Clearly, $\mu \in O$ if and only if for every countable suitable model \mathcal{M} such that $\mu \in \mathcal{M}$ there is $\alpha < \omega_2^{\mathcal{M}}$ such that $S_{\alpha+m}$ is nonstationary in $L[r(\mu)]^{\mathcal{M}}$ for every $m \in \Delta(r(\mu))$. Thus our family O has indeed a Π_2^1 definition. Just as in the proof of Theorem 1, to obtain a Π_2^1 -definable m.o. family in $L^{\mathbb{P}_{\omega_3}}$ consider the union of O with the set of all point measures.

Since for every real $a \in L^{\mathbb{P}_{\omega_3}}$ there is a random real over L , by Proposition 1 in $L^{\mathbb{P}_{\omega_3}}$ there are no Σ_2^1 m.o. families. The bounding number \mathfrak{b} remains ω_1 in $L^{\mathbb{P}_{\omega_3}}$, since the countable support iteration of S -proper ${}^\omega\omega$ -bounding posets is ${}^\omega\omega$ -bounding (see [1, Lemma 18] or [5]). \square

Remark 4.1. In [3] the following question was raised:

Question 1. If there is a Π_1^1 m.o. family, are all reals constructible?

This is to our knowledge still unsolved. Törnquist has recently shown that the existence of a Σ_2^1 m.o. family implies the existence of a Π_1^1 m.o. family, and that the existence of Σ_2^1 mad family implies the existence of a Π_1^1 mad family.

References

- [1] V. Fischer, S. D. Friedman *Cardinal characteristics and projective wellorders*, Annals of Pure and Applied Logic 161 (2010), 916-922.
- [2] V. Fischer, S. D. Friedman, L. Zdomskyy *Projective wellorders and mad families with large continuum*, to appear, Annals of Pure and Applied Logic.
- [3] V. Fischer, A. Törnquist *A co-analytic maximal set of orthogonal measures*, Journal of Symbolic Logic, 75, 4, 1403 - 1414.
- [4] S. D. Friedman, L. Zdomskyy *Projective mad families*, Annals of Pure and Applied Logic 161 (2010), 1581-1587.
- [5] M. Goldstern *A taste of proper forcing*, Set theory (Curaçao, 1995; Barcelona, 1996), 71–82, Kluwer Acad. Publ., Dordrecht, 1998.
- [6] J. I. Ihoda, S. Shelah Δ_2^1 -sets of reals, Ann. Pure Appl. Logic, 42 (1989), no. 3, 201-223. MR 998607(90f:03081).
- [7] T. Jech *Set theory*, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003.
- [8] B. Kastermans, J. Steprāns, Y. Zhang *Analytic and coanalytic families of almost disjoint functions*, Journal of Symbolic Logic 73 (2008), 1158-1172.
- [9] A. S. Kechris *Classical Descriptive Set Theory* Graduate Texts in Mathematics, 156, Springer-Verlag, 1995.
- [10] A. Kechris, N. E. Sofronidis *A strong ergodic property of unary and self-adjoint operators* Ergodic Theory and Dynamical Systems 21 (2001).
- [11] A. Miller *Infinite combinatorics and definability*, Annals of Pure and Applied Logic, vol. 41 (1989), 179-203.
- [12] A.R.D. Mathias *Happy Families*, Annals of Mathematical Logic 12 (1977), 59-111.
- [13] D. Preiss, J. Rataj *Maximal Sets of Orthogonal Measures are not Analytic* Proc. of the Amer. Math. Soc. 93, No. 3 (1985), 471-476.
- [14] D. Raghavan *Maximal almost disjoint families of functions*, Fundamenta Mathematicae 204 (2009), 241-282.