

LARGE CARDINALS AND DEFINABLE WELL-ORDERS, WITHOUT THE GCH

SY-DAVID FRIEDMAN AND PHILIPP LÜCKE

ABSTRACT. We show that there is a class-sized partial order \mathbb{P} with the property that forcing with \mathbb{P} preserves ZFC, supercompact cardinals, inaccessible cardinals and the value of 2^κ for every inaccessible cardinal κ and, if κ is an inaccessible cardinal and A is an arbitrary subset of ${}^\kappa\kappa$, then there is a \mathbb{P} -generic extension of the ground model V in which A is definable in $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters.

We use this result to construct a class-sized partial order with the above preservation properties that forces the existence of well-orders of $H(\kappa^+)$ definable in the structure $\langle H(\kappa^+), \in \rangle$ for every inaccessible cardinal κ . Assuming GCH, David Asperó and Sy-David Friedman showed in [AF09] and [AF] that there is a class-sized partial order preserving ZFC and various large cardinals and forcing the existence of a well-order of the universe whose restriction to $H(\kappa^+)$ is definable in $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a parameter-free formula for every uncountable regular cardinal κ . Our second result can be interpreted as a boldface version of this result in the absence of the GCH.

1. INTRODUCTION

Given an uncountable regular cardinal κ , we call the set ${}^\kappa\kappa$ consisting of all functions $f : \kappa \rightarrow \kappa$ the *generalized Baire Space for κ* . The study of the *descriptive set theory* of these spaces, i.e. of their definable subsets and the structural properties of these subsets, was initiated by Alan Mekler and Jouko Väänänen in [MV93] and deep links to model theory and logic were established (see [Vää95], [TV99], [Vää11] and [FHK]). A discussion of some of these results is contained in Chapter IV of [FHK]. In this paper, we study the definable subsets of this space when κ is a large cardinal, especially a supercompact cardinal.

Remember that an uncountable cardinal κ is γ -*supercompact* with $\gamma \geq \kappa$ if there is an elementary embedding $j : V \rightarrow M$ with $\text{crit}(j) = \kappa$, $\gamma < j(\kappa)$ and ${}^\gamma M \subseteq M$. This is equivalent to the existence of a normal ultrafilter on the set $\mathcal{P}_\kappa(\gamma)$ of all subsets of γ of cardinality less than κ (see [Kan03, Theorem 22.7]). Given such an ultrafilter U , we let M_U denote the transitive collapse of the corresponding ultrapower $\text{Ult}_U(V)$ and $j_U : V \rightarrow M_U$ denote the corresponding elementary embedding. Finally, we call a cardinal κ *supercompact* if κ is γ -supercompact for all $\gamma \geq \kappa$.

2000 *Mathematics Subject Classification.* 03E35, 03E47, 03E55.

Key words and phrases. Definability, definable well-orders, supercompactness, large cardinal preservation.

The first author would like to thank the FWF (Austrian Science Fund) for its support through grant #P 22430-N13.

The second author would like to thank the Deutsche Forschungsgemeinschaft for its support through grant SCHI 484/4-1 and SFB 878.

Let κ be a supercompact cardinal and A be an arbitrary subset of ${}^\kappa\kappa$. We want to construct an outer model W of the ground model V such that κ is still supercompact in W , $(2^\kappa)^V = (2^\kappa)^W$ and A is definable in the structure $\langle H(\kappa^+)^W, \in \rangle$. By extending coding methods developed in [Lüc], this aim is achieved in the following theorem.

Theorem 1.1. *There is a ZFC-preserving class forcing \mathbb{P} definable without parameters that satisfies the following statements.*

- (i) *Let κ be a cardinal with the property that there is no singular limit of inaccessible cardinals ν with $\nu^+ < \kappa \leq 2^\nu$. Then forcing with \mathbb{P} does not collapse κ and, if κ is regular, then \mathbb{P} preserves the regularity of κ .*
- (ii) *\mathbb{P} preserves the inaccessibility of inaccessible cardinals and the supercompactness of supercompact cardinals.*
- (iii) *If α is an inaccessible cardinal and G is \mathbb{P} generic over V , then $(2^\alpha)^V = (2^\alpha)^{V[G]}$.*
- (iv) *If κ is an inaccessible cardinal and A is a subset of ${}^\kappa\kappa$, then there is a condition p in \mathbb{P} with the property that A is definable in $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters whenever G is \mathbb{P} -generic over V with $p \in G$.*

In addition, if the class of inaccessible cardinals is bounded in On , then \mathbb{P} is forcing equivalent to a set-sized forcing.

In particular, if the *Singular Cardinal Hypothesis* holds in the ground model, then forcing with \mathbb{P} preserves cofinalities and cardinalities.

The proof of this result will actually show that certain degrees of supercompactness are preserved. Let κ be γ -supercompact such that γ is a cardinal with $\gamma = \gamma^{<\kappa}$, $2^\gamma = \gamma^+$ and $2^\nu \leq \gamma$, where ν is the supremum of all inaccessible cardinals smaller or equal to γ . Then κ will still be γ -supercompact after forcing with \mathbb{P} . Given a supercompact κ , we will use a classical result due to Robert Solovay to show that there is a proper class of cardinals γ that satisfy the above properties with respect to κ .

We want to use the above coding result to produce ZFC-models with definable well-orders of $H(\kappa^+)$ for every supercompact cardinal κ . We give a brief overview of related existing results. A detailed discussion of this topic can be found in the first part of [Fri10]. In [FH_a], Peter Holy and the first author constructed a class forcing that adds such definable well-orders of low quantifier complexity and preserves various large cardinals.

Theorem 1.2 ([Fri10, Theorem 9]). *There is a class forcing which forces GCH, preserves all supercompact cardinals (as well as a proper class of n -huge cardinals for each $n < \omega$) and adds a well-order of $H(\kappa^+)$ that is definable in $\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameters for every uncountable regular cardinal κ .*

If the GCH holds in the ground model, then results due to David Asperó and the first author show that it is possible to produce *lightface* definable well-orders of $H(\kappa^+)$ for every uncountable regular cardinal κ .

Theorem 1.3 ([AF09, Theorem 1.1] and [AF, Theorem 1.1]). *Assume GCH. There is a formula $\varphi(x, y)$ without parameters and a definable class-sized partial order \mathbb{P} preserving ZFC, GCH and cofinalities that satisfy the following statements.*

- (i) *\mathbb{P} forces that there is a well-order \leq of the universe such that*

$$\{\langle a, b \rangle \in H(\kappa^+)^2 \mid \langle H(\kappa^+), \in \rangle \models \varphi(a, b)\}$$

is the restriction $\leq \upharpoonright H(\kappa^+)$ and is a well-order of $H(\kappa^+)$ whenever κ is a regular uncountable cardinal.

- (ii) For all regular cardinals $\kappa \leq \lambda$, if κ is a λ -supercompact cardinal in V , then κ remains λ -supercompact after forcing with \mathbb{P} .

The second result of this paper shows that it is possible to add definable well-orders of $H(\kappa^+)$ for every inaccessible cardinal κ without assuming GCH with a class forcing that preserves supercompact cardinals and failures of the GCH at inaccessible cardinals.

Theorem 1.4. *There is a ZFC-preserving class forcing \mathbb{P} definable without parameters that satisfies the following statements.*

- (i) Let κ be a cardinal with the property that there is no singular limit of inaccessible cardinals ν with $\nu^+ < \kappa \leq 2^\nu$. Then forcing with \mathbb{P} does not collapse κ and, if κ is regular, then \mathbb{P} preserves the regularity of κ .
- (ii) \mathbb{P} preserves the inaccessibility of inaccessible cardinals and the supercompactness of supercompact cardinals.
- (iii) If α is an inaccessible cardinal and G is \mathbb{P} generic over V , then $(2^\alpha)^V = (2^\alpha)^{V[G]}$ and there is a well-order of $H(\kappa^+)^{V[G]}$ that is definable in the structure $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a formula with parameters.

In fact, the partial order \mathbb{P} constructed in the proof of this result satisfies the statements listed in Theorem 1.1.

2. GENERIC TREE CODING

The goal of this section is to construct a partial order that forces an arbitrary subset A of ${}^\kappa\kappa$ to be definable in $\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameters. This construction will be a variation of the *generic tree coding* developed in [Lüc]. In this section, we present a detailed discussion of the properties of this forcing verified in [Lüc], because most of these results will be needed in later proofs. In order to define this partial order, we give a brief review of our notation.

Given an ordinal λ and a set X , we let ${}^{<\lambda}X$ denote the set of all functions f with $\text{dom}(f) \in \lambda$ and $\text{ran}(f) \subseteq X$. If κ is a cardinal, then we let $\kappa^{<\lambda}$ denote the cardinality of ${}^{<\lambda}\kappa$. We call a set $T \subseteq ({}^{<\lambda}X)^n$ a *subtree of $({}^{<\lambda}X)^n$* if the following statements hold.

- (i) For all $\langle s_0, \dots, s_{n-1} \rangle \in T$, $\text{lh}(s_0) = \dots = \text{lh}(s_{n-1})$.
- (ii) If $\langle s_0, \dots, s_{n-1} \rangle \in T$ and $\alpha < \text{lh}(s_0)$, then $\langle s_0 \upharpoonright \alpha, \dots, s_{n-1} \upharpoonright \alpha \rangle \in T$.

Given $t = \langle t_0, \dots, t_{n-1} \rangle \in T$, we define $\text{lh}(t) = \text{lh}(t_0)$ and call the ordinal $\text{ht}(T) = \text{lub}\{\text{lh}(t) \mid t \in T\}$ the *height of T* . A tuple of functions $\langle x_0, \dots, x_{n-1} \rangle \in ({}^{\text{ht}(T)}X)^n$ is called a *cofinal branch through T* if $\langle x_0 \upharpoonright \alpha, \dots, x_{n-1} \upharpoonright \alpha \rangle \in T$ for all $\alpha < \text{ht}(T)$. We let $[T]$ denote the set of all cofinal branches through T . If T is a subtree of $({}^{<\lambda}X)^{n+1}$ for some $\lambda \in \text{On}$, then we define

$$p[T] = \{\langle x_0, \dots, x_{n-1} \rangle \in ({}^{\text{ht}(T)}X)^n \mid (\exists x_n) \langle x_0, \dots, x_n \rangle \in [T]\}.$$

Definition 2.1. Let κ be an infinite cardinal. A subset A of ${}^\kappa\kappa$ is a Σ_1^1 -subset if there is a subtree T of ${}^{<\kappa}\kappa \times {}^{<\kappa}\kappa$ with $A = p[T]$.

Given an uncountable regular cardinal κ with $\kappa = \kappa^{<\kappa}$, it is a well-known fact that a subset of ${}^\kappa\kappa$ is a Σ_1^1 -subset if and only if it is definable in the structure

$\langle H(\kappa^+), \in \rangle$ by a Σ_1 -formula with parameters. A proof of this folklore result can be found in [Lüc, Section 2].

We sketch the idea behind the definition of our forcing notion. Fix an uncountable regular cardinal κ with $\kappa = \kappa^{<\kappa}$ and an enumeration $\langle s_\alpha \mid \alpha < \kappa \rangle$ of all elements in ${}^{<\kappa}\kappa$. We say that $x \in {}^\kappa\kappa$ is *coded by* $z \in {}^{<\kappa}2$ and $\gamma < \kappa$ if

$$s_\beta \subseteq x \iff z(\langle \gamma, \beta \rangle) = 1$$

holds for all $\beta < \kappa$, where $\langle \cdot, \cdot \rangle$ denote the Gödel-pairing function. Given a subset A of ${}^\kappa\kappa$, our forcing will add a subtree T_G of ${}^{<\kappa}2$ with the property that, in the generic extension, A is equal to the set of all x that are coded by some $z \in [T_G]$ and $\gamma < \kappa$. This definition of A provides a tree T in the generic extension that satisfies $A = p[T]$.

Definition 2.2. Given a limit ordinal λ , we call a pair $\langle A, s \rangle$ a λ -*coding basis* if the following statements hold.

- (i) A is a non-empty subset of ${}^\lambda\lambda$ and $s : \lambda \rightarrow {}^{<\lambda}\lambda$.
- (ii) $\text{ran}(s)$ contains $\{x \upharpoonright \alpha \mid x \in A, \alpha < \lambda\}$ and all constant functions in ${}^{<\lambda}\lambda$.
- (iii) For all $\alpha < \lambda$, $\text{lh}(s(\alpha)) \leq \alpha$ and $\{\beta < \lambda \mid s(\alpha) = s(\beta)\}$ is unbounded in λ .

For the rest of this section, we fix a regular uncountable cardinal κ with $\kappa = \kappa^{<\kappa}$. Given a κ -coding basis $\langle A, s \rangle$, we define a partial order $\mathbb{P}_s(A)$. The domain of $\mathbb{P}_s(A)$ consists of all triples $p = \langle T_p, f_p, h_p \rangle$ with the following properties.

- (i) T_p is a subtree of ${}^{<\kappa}2$ that satisfies the following statements.
 - (a) T_p has cardinality less than κ .
 - (b) If $t \in T_p$ with $\text{lh}(t) + 1 < \text{ht}(T_p)$, then t has two immediate successors in T_p .
- (ii) $f_p : A \xrightarrow{\text{part}} [T_p]$ is a partial function such that $\text{dom}(f_p)$ is a non-empty set of cardinality less than κ .
- (iii) $h_p : A \xrightarrow{\text{part}} \kappa$ is a partial function with the following properties.
 - (a) $\text{dom}(h_p) = \text{dom}(f_p)$.
 - (b) For all $x \in \text{dom}(h_p)$ and $\alpha, \beta < \text{ht}(T_p)$ with $\alpha = \langle h_p(x), \beta \rangle$, we have

$$s(\beta) \subseteq x \iff f_p(x)(\alpha) = 1.$$

We define $p \leq_{\mathbb{P}_s(A)} q$ to hold if following statements are satisfied.

- (a) T_p is either equal to T_q or an end-extension of T_q .
- (b) If $x \in \text{dom}(f_q)$, then $x \in \text{dom}(f_p)$ and $f_q(x)$ is an initial segment of $f_p(x)$.
- (c) $h_q = h_p \upharpoonright \text{dom}(h_q)$.

Lemma 2.3. $\mathbb{P}_s(A)$ is $<\kappa$ -closed, satisfies the κ^+ -chain condition and has cardinality at most 2^κ .

Proof. If $\lambda \in \text{Lim} \cap \kappa$ and $\langle p_\mu \mid \mu < \lambda \rangle$ is a strictly $\leq_{\mathbb{P}_s(A)}$ -descending sequence in $\mathbb{P}_s(A)$, then we define $T = \bigcup_{\mu < \lambda} T_{p_\mu}$, $h = \bigcup_{\mu < \lambda} h_\mu$ and

$$f(x) = \bigcup \{f_{p_\mu}(x) \mid \mu < \lambda, x \in \text{dom}(f_{p_\mu})\}$$

for all $x \in \text{dom}(h)$. It is easy to see that $p = \langle T, f, h \rangle \in \mathbb{P}_s(A)$ and $p \leq_{\mathbb{P}_s(A)} p_\mu$ holds for all $\mu < \lambda$.

Next, assume that $\langle p_\mu \mid \mu < \kappa^+ \rangle$ enumerates an antichain in $\mathbb{P}_s(A)$. By our assumptions, we can assume $T_{p_\mu} = T_{p_\rho}$ for all $\mu, \rho < \kappa^+$. A Δ -system argument shows that we may assume the existence of an $r \subseteq A$ with $r = \text{dom}(f_{p_\mu}) \cap \text{dom}(f_{p_\rho})$,

$f_{p_\mu} \upharpoonright r = f_{p_\rho} \upharpoonright r$ and $h_{p_\mu} \upharpoonright r = h_{p_\rho} \upharpoonright r$ for all $\mu < \rho < \kappa^+$. But this shows that $\langle T_{p_0}, f_{p_0} \cup f_{p_1}, h_{p_0} \cup h_{p_1} \rangle$ is a common extension of p_0 and p_1 , a contradiction.

Finally, the assumption $\kappa = \kappa^{<\kappa}$ implies that there are only κ -many such subtrees and 2^κ -many such partial functions of cardinality less than κ . \square

The next lemma will allow us to show that various subsets of $\mathbb{P}_s(A)$ are dense.

Lemma 2.4. *Fix a condition p in $\mathbb{P}_s(A)$ and a sequence $\langle c_x \in {}^\kappa 2 \mid x \in \text{dom}(f_p) \rangle$. There exists a $\leq_{\mathbb{P}_s(A)}$ -descending sequence $\langle p_\mu \in \mathbb{P}_s(A) \mid \text{ht}(T_{p_\mu}) \leq \mu < \kappa \rangle$ such that $p = p_{\text{ht}(T_p)}$ and the following statements hold for all $\text{ht}(T_{p_\mu}) \leq \mu < \kappa$.*

- (i) $\text{dom}(f_{p_\mu}) = \text{dom}(f_p)$ and $\text{ht}(T_{p_\mu}) = \mu$.
- (ii) If $x \in \text{dom}(f_p)$ and $\mu \neq \langle h_p(x), \beta \rangle$ for all $\beta < \kappa$, then

$$f_{p_{\mu+1}}(x)(\mu) = c_x(\mu).$$

- (iii) If $\mu \in \text{Lim}$, then $\text{ran}(f_{p_\mu}) = T_{p_{\mu+1}} \cap {}^\mu 2$.

Proof. We construct the sequences inductively. If $\mu \in \text{Lim}$, then we define $T_{p_\mu} = \bigcup \{T_{p_{\bar{\mu}}} \mid \text{ht}(T_{p_{\bar{\mu}}}) \leq \bar{\mu} < \mu\}$. Given $x \in \text{dom}(f_p)$, we define

$$f_{p_\mu}(x) = \bigcup \{f_{p_{\bar{\mu}}}(x) \mid \text{ht}(T_{p_{\bar{\mu}}}) \leq \bar{\mu} < \mu\}.$$

If $\mu = \bar{\mu} + 1$ with $\bar{\mu} \notin \text{Lim}$, then $T_{p_{\bar{\mu}}}$ has a maximal level and there is only one suitable tree T_{p_μ} of height μ end-extending it. In particular, $f_{p_{\bar{\mu}}}(x) \in T_{p_\mu}$ for all $x \in \text{dom}(f_p)$. For all $x \in \text{dom}(f_p)$, we define $f_{p_\mu}(x)$ to be the unique element t of ${}^\mu 2$ with $f_{p_{\bar{\mu}}}(x) \subseteq t$ and

$$t(\bar{\mu}) = \begin{cases} 1, & \text{if } \bar{\mu} = \langle h_p(x), \beta \rangle \text{ and } s(\beta) \subseteq x, \\ 0, & \text{if } \bar{\mu} = \langle h_p(x), \beta \rangle \text{ and } s(\beta) \not\subseteq x, \\ c_x(\bar{\mu}), & \text{otherwise.} \end{cases}$$

Finally, if $\mu = \bar{\mu} + 1$ with $\bar{\mu} \in \text{Lim}$, then we set $T_{p_\mu} = T_{p_{\bar{\mu}}} \cup \text{ran}(f_{p_{\bar{\mu}}})$ and define f_{p_μ} as in the first successor case. \square

Corollary 2.5. *The following sets are dense subsets of $\mathbb{P}_s(A)$.*

- (i) $C_\mu = \{p \in \mathbb{P}_s(A) \mid \text{ht}(T_p) > \mu\}$ for all $\mu < \kappa$.
- (ii) $D_x = \{p \in \mathbb{P}_s(A) \mid x \in \text{dom}(f_p)\}$ for all $x \in A$.
- (iii) $E_{x,y} = \{p \in \mathbb{P}_s(A) \mid x, y \in \text{dom}(f_p), f_p(x) \neq f_p(y)\}$ for all $x, y \in A$.
- (iv) $F_z = \{p \in \mathbb{P}_s(A) \mid \text{ht}(T_p) = \mu + 1, z \upharpoonright \mu \notin T_p\}$ for all $z \in {}^\kappa 2$.

Proof. (i) This statement follows directly from Lemma 2.4.

- (ii) Given $p \in \mathbb{P}_s(A)$ with $x \notin \text{dom}(f_p)$ and $b \in [T_p]$, we define

$$q = \langle T_p, f_p \cup \{\langle x, b \rangle\}, h_p \cup \{\langle x, \text{ht}(T_p) \rangle\} \rangle.$$

Then $q \in D_x$ and $q \leq_{\mathbb{P}_s(A)} p$.

(iii) Given $p \in \mathbb{P}_s(A)$, we can apply the above result to find $q \leq_{\mathbb{P}_s(A)} p$ with $x, y \in \text{dom}(f_q)$. There is $\text{ht}(T_q) \leq \mu < \kappa$ with $\langle h_q(x), \beta_0 \rangle \neq \mu \neq \langle h_q(y), \beta_1 \rangle$ for all $\beta_0, \beta_1 < \kappa$ and we can use Lemma 2.4 to find $q^* \leq_{\mathbb{P}_s(A)} q$ with $\text{ht}(T_{q^*}) = \mu + 1$ and $f_{q^*}(x)(\mu) \neq f_{q^*}(y)(\mu)$.

(iv) Fix $p \in \mathbb{P}_s(A)$ and $\text{ht}(T_p) \leq \mu < \kappa$ with $\mu \neq \langle h_p(x), \beta \rangle$ for all $x \in \text{dom}(f_p)$ and $\beta < \kappa$. Using Lemma 2.4, we can find $q \leq_{\mathbb{P}_s(A)} p$ with $\text{ht}(T_q) = \mu + 1$, $\text{dom}(f_q) = \text{dom}(f_p)$ and $f_q(x)(\mu) = 1 - z(\mu)$ for all $x \in \text{dom}(f_p)$. In particular, $z \upharpoonright (\mu + 1) \notin \text{ran}(f_q)$. Another application of the above lemma gives us conditions $s \leq_{\mathbb{P}_s(A)} r \leq_{\mathbb{P}_s(A)} q$ with $\text{ht}(T_s) = \text{ht}(T_r) + 1 = \text{ht}(T_q) + \omega + 1$, $\text{dom}(f_s) = \text{dom}(f_p)$

and $T_s \cap \text{ht}(T_r)2 = \text{ran}(f_r)$. Since $z \upharpoonright \text{ht}(T_r) \neq f_r(x)$ for all $x \in \text{dom}(f_p)$, we have $z \upharpoonright \text{ht}(T_r) \notin T_s$. \square

Corollary 2.6. *Let G be $\mathbb{P}_s(A)$ -generic over V . The following statements hold true in $V[G]$.*

- (i) $T_G = \bigcup_{p \in G} T_p$ is a subtree of ${}^{<\kappa}2$ of height κ with $[T_G] \cap V = \emptyset$.
- (ii) If we define $F_G(x) = \bigcup \{f_p(x) \mid p \in G, x \in \text{dom}(f_p)\}$ for all $x \in A$, then $F_G : A \rightarrow [T_G]$ is an injection.
- (iii) Let $H_G = \bigcup_{p \in G} h_p$. Then $H_G : A \rightarrow \kappa$ and

$$(1) \quad s(\beta) \subseteq x \iff F_G(x)(\prec H_G(x), \beta \succ) = 1$$

for all $x \in A$ and $\beta < \kappa$. \square

Lemma 2.7. *If G be $\mathbb{P}_s(A)$ -generic over V , then $\text{ran}(F_G) = [T_G]^{V[G]}$.*

Proof. Let $\dot{T} \in V^{\mathbb{P}_s(A)}$ be the canonical name for T_G and $\dot{F} \in V^{\mathbb{P}_s(A)}$ be the canonical name for F_G .

Assume, toward a contradiction, that there is an $x \in [T_G]^{V[G]} \setminus \text{ran}(F_G)$ and let $\tau \in V^{\mathbb{P}_s(A)}$ be a name for x . By the above corollary, $x \notin V$ and there is a $p_0 \in G$ with

$$p_0 \Vdash \text{“}\tau \in [\dot{T}] \wedge \tau \notin \check{V} \wedge \tau \notin \text{ran}(\dot{F})\text{”}.$$

For each $r \leq_{\mathbb{P}_s(A)} p$, we define a partial function $t_r : \kappa \xrightarrow{\text{part}} 2$ in V by setting

$$t_r = \bigcup \{t \in {}^{<\kappa}2 \mid r \Vdash \text{“}\check{t} \subseteq \tau\text{”}\}.$$

We have $t_r \in {}^{<\kappa}2$, because $r \Vdash \text{“}\tau \notin \check{V}\text{”}$. Note that $r_1 \leq_{\mathbb{P}_s(A)} r_0 \leq_{\mathbb{P}_s(A)} p$ implies $t_{r_0} \subseteq t_{r_1}$. Since $\mathbb{1}_{\mathbb{P}_s(A)} \Vdash \text{“}\tau \upharpoonright \check{\alpha} \in \check{V}\text{”}$ holds for all $\alpha < \kappa$, the set $\{r \leq_{\mathbb{P}_s(A)} p \mid \alpha \subseteq \text{dom}(t_r)\}$ is dense below p for all $\alpha < \kappa$.

Moreover, if $p' \leq_{\mathbb{P}_s(A)} p$, then we can find a $p'' \leq_{\mathbb{P}_s(A)} p'$ with the property that for every $x \in \text{dom}(f_{p'})$ there is an $\alpha < \text{ht}(T_{p''}) \cap \text{dom}(t_{p''})$ with $f_{p''}(x)(\alpha) \neq t_r(\alpha)$, because $\mathbb{P}_s(A)$ is $<\kappa$ -closed.

Given $p_0 \leq_{\mathbb{P}_s(A)} p$, the above remarks allow us to construct a strictly $\leq_{\mathbb{P}_s(A)}$ -descending sequence $\langle p_n \in \mathbb{P}_s(A) \mid n < \omega \rangle$ with the following properties.

- (i) For all $n < \omega$, $\text{ht}(T_{p_n}) \subseteq \text{dom}(t_{p_{n+1}})$ and $\text{dom}(t_{p_n}) \subsetneq \text{ht}(T_{p_{n+1}})$.
- (ii) For all $n < \omega$ and $x \in \text{dom}(f_{p_n})$, there is an $\alpha \in \text{dom}(t_{p_{n+1}})$ with

$$f_{p_{n+1}}(x)(\alpha) \neq t_{p_{n+1}}(\alpha).$$

The proof of Lemma 2.3 shows that there exists a greatest lower bound $p_\omega \in \mathbb{P}_s(A)$ of the sequence $\langle p_n \mid n < \omega \rangle$. This means $T_{p_\omega} = \bigcup_{n < \omega} T_{p_n}$ and $\text{dom}(f_{p_\omega}) = \bigcup_{n < \omega} \text{dom}(f_{p_n})$. Let $t = t_{p_\omega} \upharpoonright \text{ht}(T_{p_\omega})$. Since $p_\omega \Vdash \text{“}\check{t} \subseteq \tau \wedge \tau \in [\dot{T}]\text{”}$, we have $p_\omega \Vdash \text{“}\check{t} \in \dot{T}\text{”}$.

By our construction, we have $\mu = \text{ht}(T_{p_\omega}) \in \text{Lim}$ and $t \notin \text{ran}(f_{p_\omega})$. We can apply Lemma 2.4 to find a condition $p^* \leq_{\mathbb{P}_s(A)} p_\omega$ with $\text{ht}(T_{p^*}) = \mu + 1$ and $t \notin T_{p^*}$. This obviously implies $p^* \Vdash \text{“}\check{t} \notin \dot{T}\text{”}$, a contradiction. \square

Lemma 2.8. *Let G be $\mathbb{P}_s(A)$ -generic over V . The following statements are equivalent for $y \in ({}^\kappa \kappa)^{V[G]}$.*

- (i) $y \in A$.

(ii) *There is $z \in [T_G]^{V[G]}$ and $\gamma < \kappa$ such that*

$$(2) \quad s(\beta) \subseteq y \iff z(\prec\gamma, \beta\succ) = 1$$

holds for all $\beta < \kappa$.

Proof. If $y \in A$, then Corollary 2.6 shows that $F_G(y) \in [T_G]$ and $H_G(y) < \kappa$ witness that the second statement holds true.

Pick $y \in (\kappa^\kappa)^{V[G]}$, $z \in [T_G]^{V[G]}$ and $\gamma < \kappa$ such that (2) holds. By Lemma 2.7, we have $z = F_G(x)$ for some $x \in A$. Pick $p \in G$ with $x \in \text{dom}(f_p)$. Assume, toward a contradiction, that $\gamma \neq h_p(x) = H_G(x)$. By Lemma 2.4 and our assumptions on s , this implies that the set

$$D_t = \{q \leq_{\mathbb{P}_s(A)} p \mid \text{ht}(T_q) = \mu + 1, \mu = \prec\gamma, \beta\succ, f_q(x)(\mu) = 0, s(\beta) = t\}$$

is dense below p for all $t \in \text{ran}(s)$ and there is a $q \in G \cap D_{y \upharpoonright 1}$ with $q \leq_{\mathbb{P}_s(A)} p$. Then there is a $\beta < \kappa$ with $\text{ht}(T_q) = \prec\gamma, \beta\succ + 1$, $z(\prec\gamma, \beta\succ) = 0$ and $s(\beta) = y \upharpoonright 1 \subseteq y$, contradicting (2). This shows $\gamma = H_G(x)$ and we can conclude that

$$s(\beta) \subseteq y \iff z(\prec\gamma, \beta\succ) = 1 \iff F_G(x)(\prec H_G(x), \beta\succ) = 1 \iff s(\beta) \subseteq x$$

holds for all $\beta < \kappa$. Since every initial segment of x is of the form $s(\beta)$ for some $\beta < \kappa$, we can conclude $y = x \in A$. \square

Theorem 2.9 ([Lüc, Theorem 1.5]). *If G is $\mathbb{P}_s(A)$ -generic over V , then A is a Σ_1^1 -subset of ${}^\kappa\kappa$ in $V[G]$.*

Proof. In $V[G]$, define T to be the set that consists of pairs $\langle t, u \rangle$ such that $t \in {}^{<\kappa}\kappa$, $u \in {}^{<\kappa}\kappa$ and there is $\gamma < \kappa$ and $v \in T_G$ with $\text{lh}(z) = \text{lh}(u) = \text{lh}(v)$, $u(\alpha) = \prec\gamma, v(\alpha)\succ$ for all $\alpha < \text{lh}(s)$ and

$$s(\beta) \subseteq t \iff v(\prec\gamma, \beta\succ) = 1$$

for all $\beta < \text{lh}(s)$ with $\prec\gamma, \beta\succ < \text{lh}(s)$. It is easy to check that T is a tree.

If $\langle x, y \rangle \in [T]^{V[G]}$, then there is $z \in [T_G]^{V[G]}$ and $\gamma < \kappa$ with $y(\beta) = \prec\gamma, z(\beta)\succ$ and

$$s(\beta) \subseteq x \iff z(\prec\gamma, \beta\succ) = 1$$

for all $\beta < \kappa$. By Lemma 2.8, this implies $x \in A$.

Conversely, if $x \in A$ and $y \in (\kappa^\kappa)^{V[G]}$ with $y(\alpha) = \prec H_G(x), F_G(x)(\alpha)\succ$, then $\langle x, y \rangle \in [T]$ by our assumptions on s and Lemma 2.8. \square

We close this section by proving a structural property of our coding forcing that will be needed in the proof of supercompactness preservation.

Lemma 2.10. *Assume $P \subseteq \mathbb{P}_s(A)$ satisfies the following properties.*

- (i) $\eta = \text{lub}\{\text{ht}(T_p) \mid p \in P\} \in \text{Lim} \cap \kappa$.
- (ii) $D = \bigcup\{\text{dom}(f_p) \mid p \in P\}$ has cardinality less than κ .
- (iii) If $p_0, p_1 \in P$, then there is $q \in P$ with $q \leq_{\mathbb{P}_s(A)} p_0, p_1$.

Then there is a unique condition $p_P \in \mathbb{P}_s(A)$ with $\text{ht}(T_{p_P}) = \eta$, $\text{dom}(f_{p_P}) = D$ and $p_P \leq_{\mathbb{P}_s(A)} p$ for all $p \in P$.

Proof. Set $T = \bigcup\{T_p \mid p \in P\}$. Then T is a tree of height η and an end-extension of T_p for all $p \in P$. If we define

$$F : D \longrightarrow [T]; x \longmapsto \bigcup\{f_p(x) \mid p \in P, x \in \text{dom}(f_p)\} \in [T],$$

then this is a well-defined function. Moreover, for all $x \in D$ there is a unique $H(x) < \kappa$ with $h_p(x) = H(x)$ for all $p \in P$ with $x \in \text{dom}(f_p)$ and we can define $H : D \rightarrow \kappa$ in this way.

If $x \in D$ and $\alpha, \beta < \eta$ with $\alpha = \prec H(x), \beta \succ$, then there is $p \in P$ with $x \in \text{dom}(f_p)$ and $\alpha, \beta < \text{ht}(T_p)$. We can conclude

$$s(\beta) \subseteq x \iff f_p(x)(\alpha) = 1 \iff F(x)(\alpha) = 1.$$

This shows that $p_P = \langle T, F, H \rangle$ is a condition in \mathbb{P} with $p_P \leq p$ for all $p \in P$.

Let $q \in \mathbb{P}_s(A)$ be a condition with $\text{ht}(T_q) = \eta$, $\text{dom}(f_q) = D$ and $q \leq_{\mathbb{P}_s(A)} p$ for all $p \in P$. Since $\eta \in \text{Lim}$, for every $t \in T_q$ there is a $p \in P$ with $\text{lh}(t) < \text{ht}(T_p)$ and therefore $t \in T_p$. This shows $T_q = \bigcup \{T_p \mid p \in P\} = T$. In the same way, we can show $f_q(x) = \bigcup \{f_p(x) \mid p \in P, x \in \text{dom}(f_p)\} = F(x)$ and $h_q(x) = H(x)$ for all $x \in D$. This means $q = p_P$. \square

3. CODING WELL-ORDERS

In this section, we show how to apply the results of the last section to construct a definable well-order of $\mathbb{H}(\kappa^+)$ in a $\mathbb{P}_s(A)$ -generic extension of the ground model. Throughout this section κ is an uncountable regular cardinal with $\kappa = \kappa^{<\kappa}$.

Given functions $x, y \in {}^\kappa\kappa$, let $\prec x, y \succ$ denote the unique function $z \in {}^\kappa\kappa$ such that

$$z(\prec \alpha, \beta \succ) = \begin{cases} x(\beta), & \text{if } \alpha = 0, \\ y(\beta), & \text{if } \alpha = 1, \\ 0, & \text{otherwise} \end{cases}$$

holds for all $\alpha, \beta < \kappa$. We say that a κ -coding basis $\langle A, s \rangle$ codes a well-order of ${}^\kappa\kappa$ if there is a well-order \leq of ${}^\kappa\kappa$ such that $A = \{\prec x, y \succ \mid x, y \in {}^\kappa\kappa, x \leq y\}$.

Theorem 3.1. *If $\langle A, s \rangle$ is a κ -coding basis that codes a well-order of ${}^\kappa\kappa$ and G is $\mathbb{P}_s(A)$ -generic over V , then there is a well-order of $\mathbb{H}(\kappa^+)$ that is definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$ by a formula with parameters.*

Proof. We work in $V[G]$. Let \leq denote the well-order of $({}^\kappa\kappa)^V$ coded by A . By Theorem 2.9, both \leq and $({}^\kappa\kappa)^V$ are definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$.

Define R to be the set of all pairs $\langle a, x \rangle$ in $\mathbb{H}(\kappa^+) \times {}^\kappa 2$ such that there is a bijection $b : \kappa \rightarrow \text{tc}(\{a\} \cup \kappa)$ with the following properties.

- (i) For all $\alpha, \beta < \kappa$, $x(\prec 0, \prec \alpha, \beta \succ \succ) = 1$ if and only if $b(\alpha) \in b(\beta)$.
- (ii) For all $\alpha < \kappa$, $x(\prec 1, \alpha \succ) = 1$ if and only if $b(\alpha) \in a$.

This relation is definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$. If $\langle a_0, x \rangle, \langle a_1, x \rangle \in R$, then it is easy to see that $a_0 = a_1$ holds. Moreover, if $\langle a, x \rangle \in R$ and $x \in V$, then a is an element of $\mathbb{H}(\kappa^+)^V$. This shows that $\mathbb{H}(\kappa^+)^V$ is definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$.

Since $\mathbb{P}_s(A)^V$ is $<\kappa$ -closed and A is definable in the above structure, we have $\mathbb{P}_s(A)^V = \mathbb{P}_s(A) \subseteq \mathbb{H}(\kappa^+)$ and this partial order is definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$. Given $y \in A$ and $\gamma < \kappa$, the proof of Lemma 2.8 shows that $H_G(y) = \gamma$ holds if and only if there is a $z \in [T_G]$ such that (2) holds for all $\beta < \kappa$. We can conclude that the function H_G is definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$. In combination with (1), this implies that the function F_G is also definable in this structure. The filter G consists of all conditions p in $\mathbb{P}_s(A)$ such that T_G is an end-extension of T_p and, if $x \in \text{dom}(f_p)$, then $f_p(x) = F_G(x) \upharpoonright \text{ht}(T_p)$ and $h_p(x) = H_G(x)$. Since all of these parameters are either elements of $\mathbb{H}(\kappa^+)$ or definable in this structure, we can conclude that G is definable in $\langle \mathbb{H}(\kappa^+), \in \rangle$.

Let N denote the set of all function $n : \kappa \times \kappa \longrightarrow \mathbb{P}_s(A)$ in V with the property that $A_\alpha^n = \{n(\alpha, \beta) \mid \beta < \kappa\}$ is an anti-chain in $\mathbb{P}_s(A)$ for all $\alpha < \kappa$. We define E to be the set consisting of all pairs $\langle y, n \rangle \in {}^\kappa 2 \times N$ such that

$$y(\alpha) = 1 \iff A_\alpha^n \cap G \neq \emptyset$$

holds for all $\alpha < \kappa$. By identifying functions in N with nice names for subsets of κ , it is easy to see that the domain of E is ${}^\kappa 2$. Both relations N and R are all definable in the above structure.

Define $r : H(\kappa^+) \longrightarrow ({}^\kappa 2)^V$ to be the function that sends $a \in H(\kappa^+)$ to the \leq -least $x \in ({}^\kappa 2)^V$ such that $R(a, y)$, $E(y, n)$ and $R(n, x)$ for some $y \in {}^\kappa 2$ and $n \in N$. This function is definable in $\langle H(\kappa^+), \in \rangle$ and yields a definable well-order of $H(\kappa^+)$. \square

Next, we introduce partial orders \mathbb{C}_α that *randomly* well-order ${}^\alpha \alpha$ if α is a regular uncountable cardinal with $\alpha = \alpha^{<\alpha}$. This coding is *random* in the sense that the generic filter chooses the well-order of ${}^\alpha \alpha$ that is coded using a partial order of the form $\mathbb{P}_s(A)$.

If α is not a regular uncountable cardinal with $\alpha = \alpha^{<\alpha}$, then we define \mathbb{C}_α to be the trivial partial order. Otherwise, we define the domain of \mathbb{C}_α to consist of conditions $\langle A, s, p \rangle$ such that either $A = s = p = \emptyset$ or $\langle A, s \rangle$ is an α -coding basis that codes a well-ordering of ${}^\alpha \alpha$ and $p \in \mathbb{P}_s(A)$. We set $\langle A, s, p \rangle \leq_{\mathbb{C}_\alpha} \langle B, t, q \rangle$ if either $B = \emptyset$ or $A = B \neq \emptyset$, $s = t$ and $p \leq_{\mathbb{P}_s(A)} q$.

Proposition 3.2. *Let α be a regular uncountable cardinal with $\alpha = \alpha^{<\alpha}$.*

- (i) \mathbb{C}_α is $<\alpha$ -closed.
 - (ii) A filter G is \mathbb{C}_α -generic over V if and only if there is an α -coding basis $\langle A, s \rangle$ coding a well-order of ${}^\alpha \alpha$ in V and $H \in \mathbb{P}_s(A)$ -generic over V with
- $$(3) \quad G = \{\langle \emptyset, \emptyset, \emptyset \rangle\} \cup \{\langle A, s, p \rangle \in \mathbb{C}_\alpha \mid p \in H\}.$$

In particular, $V[G] = V[H]$ holds in the above situation, forcing with \mathbb{C}_α preserves cofinalities, cardinalities and 2^α and every set of ordinals of cardinality at most α in a \mathbb{C}_α -generic extension of the ground model V is covered by a set that is an element of V and has cardinality α in V .

- (iii) If G is \mathbb{C}_α -generic over V , then there is a well-order of $H(\alpha^+)$ that is definable in $\langle H(\alpha^+), \in \rangle$ by a formula with parameters. \square

Note that \mathbb{C}_α is uniformly definable in parameter α .

4. ITERATED CODING FORCING

In this section, we use the coding forcing developed above in an iterated forcing construction. Our account of iterated forcing follows [Bau83] and [Cum10] and we will repeatedly use results proved there.

By the results of the last section, there is a unique forcing iteration

$$\langle \langle \vec{\mathbb{C}}_{<\alpha} \mid \alpha \in \text{On} \rangle, \langle \dot{\mathbb{C}}_\alpha \mid \alpha \in \text{On} \rangle \rangle$$

with *Easton support* (see [Cum10, Definition 7.5]) satisfying the following properties.

- (i) If $\beta < \alpha$ and α is inaccessible, then $\vec{\mathbb{C}}_{<\beta}, \dot{\mathbb{C}}_\beta \in V_\alpha$.
- (ii) If α is not an inaccessible cardinal, then $\mathbb{1}_{\vec{\mathbb{C}}_{<\alpha}} \Vdash \text{“}\dot{\mathbb{C}}_\alpha \text{ is trivial”}$.
- (iii) If α is an inaccessible cardinal, then $\mathbb{1}_{\vec{\mathbb{C}}_{<\alpha}} \Vdash \text{“}\dot{\mathbb{C}}_\alpha = \mathbb{C}_\alpha \text{”}$.

For all $\nu \leq \mu$, we let $\dot{\mathbb{C}}_{[\nu, \mu]}$ denote the canonical $\vec{\mathbb{C}}_{< \nu}$ -name with

$$\mathbb{1}_{\vec{\mathbb{C}}_{< \nu}} \Vdash \text{“}\dot{\mathbb{C}}_{[\nu, \mu]} \text{ is a partial order with domain } \{\vec{p} \upharpoonright [\check{\nu}, \check{\mu}] \mid \vec{p} \in \vec{\mathbb{C}}_{< \mu}\}\text{”}$$

such that there is a dense embedding $e_{[\nu, \mu]} : \vec{\mathbb{C}}_{< \mu} \longrightarrow \vec{\mathbb{C}}_{< \nu} * \dot{\mathbb{C}}_{[\nu, \mu]}$ with $e_{[\nu, \mu]}(\vec{p}) = \langle \vec{p} \upharpoonright \nu, \dot{q} \rangle$ and $\mathbb{1}_{\vec{\mathbb{C}}_{< \nu}} \Vdash \text{“}\dot{q} = \check{p} \upharpoonright [\check{\nu}, \check{\mu}]\text{”}$ (see [Bau83, Section 5]).

Proposition 4.1. *Let $\alpha < \mu$ and μ be a regular cardinal. Assume that there are no inaccessible cardinals in (α, μ) and $\vec{\mathbb{C}}_{< \alpha+1}$ has the property that every set of ordinals of cardinality less than μ in a $\vec{\mathbb{C}}_{< \alpha+1}$ -generic extension of the ground model is covered by a set of cardinality less than μ in the ground model. Then*

$$\mathbb{1}_{\vec{\mathbb{C}}_{< \alpha+1}} \Vdash \text{“}\dot{\mathbb{C}}_{[\alpha+1, \nu]} \text{ is } < \check{\mu}\text{-closed”}$$

for all $\nu > \alpha$.

Proof. For all $\alpha < \beta < \mu$, we have $\mathbb{1}_{\vec{\mathbb{C}}_{< \beta}} \Vdash \text{“}\dot{\mathbb{C}}_{\beta} \text{ is trivial”}$ by the definition of $\vec{\mathbb{C}}_{< \nu}$ and our assumptions on μ . This shows that $\mathbb{1}_{\vec{\mathbb{C}}_{< \beta}} \Vdash \text{“}\dot{\mathbb{C}}_{\beta} \text{ is } < \check{\mu}\text{-closed”}$ holds for all $\beta > \alpha$. Moreover, $\vec{\mathbb{C}}_{< \beta}$ is an inverse limit for every limit ordinal $\beta > \alpha$ with $\text{cof}(\beta) < \mu$. We can apply [Cum10, Proposition 7.12] to deduce the statement of the claim. \square

Proposition 4.2. *If α is an inaccessible cardinal, then $\vec{\mathbb{C}}_{< \alpha}$ preserves the inaccessibility of α .*

Proof. Let G be $\vec{\mathbb{C}}_{< \alpha}$ -generic over V . Fix $\beta < \alpha$ and let $G_{\beta+1}$ denote the corresponding filter in $\vec{\mathbb{C}}_{< \beta+1}$. If $\mu = (|\vec{\mathbb{C}}_{< \beta}|^+ + |\beta|)^+$, then there are no inaccessible cardinals in (β, μ) and $\dot{\mathbb{C}}_{[\beta+1, \alpha]}^{V[G_{\beta+1}]}$ is $< \beta^+$ -closed by Proposition 4.1. This shows $(\beta^+)^{V[G]} \subseteq V[G_{\beta+1}]$. Since $\vec{\mathbb{C}}_{< \beta+1} \in V_{\alpha}$ and α is inaccessible in $V[G_{\beta+1}]$, the statement of the claim follows directly. \square

Proposition 4.3. *$\vec{\mathbb{C}}_{< \nu}$ preserves the inaccessibility of all inaccessible cardinals.*

Proof. By Proposition 4.2 and our assumptions, $\vec{\mathbb{C}}_{< \nu}$ preserves the cofinality, cardinality and inaccessibility of all inaccessible cardinals greater or equal to ν .

Let $\alpha < \nu$ be an inaccessible cardinal. By Proposition 4.2, $\vec{\mathbb{C}}_{< \alpha}$ preserves the inaccessibility of α and $\mathbb{1}_{\vec{\mathbb{C}}_{< \alpha}} \Vdash \text{“}\dot{\mathbb{C}}_{\alpha} \text{ is not trivial”}$. Proposition 3.2 shows that $\vec{\mathbb{C}}_{< \alpha+1}$ preserves the inaccessibility of α . If $\mu = (|\vec{\mathbb{C}}_{< \alpha+1}|^+ + \alpha)^+$, then there are no inaccessible cardinals in (α, μ) and $\mathbb{1}_{\vec{\mathbb{C}}_{< \alpha+1}} \Vdash \text{“}\dot{\mathbb{C}}_{[\alpha+1, \nu]} \text{ is } < \check{\mu}\text{-closed”}$. In particular, $\vec{\mathbb{C}}_{< \nu}$ preserves the inaccessibility of α . \square

Lemma 4.4. *Let $\alpha < \nu$ and α be an inaccessible cardinal. Assume G is $\vec{\mathbb{C}}_{< \nu}$ -generic over V , \vec{G} is the corresponding filter in $\vec{\mathbb{C}}_{< \alpha}$ and G_{α} is the corresponding filter in $\dot{\mathbb{C}}_{\alpha}^{\vec{G}}$. Then $(2^{\alpha})^{V[G]} = (2^{\alpha})^V$, $\dot{\mathbb{C}}_{\alpha}^{\vec{G}} = \mathbb{C}_{\alpha}^{V[G]}$ is not the trivial partial order and, if $\langle A, s \rangle$ is an α -coding basis coding a well-order of ${}^{\alpha}\alpha$ in $V[\vec{G}]$ with $\langle A, s, \mathbb{1}_{\mathbb{P}_s(A)} \rangle \in G_{\alpha}$, then A is a Σ_1^1 -subset of ${}^{\alpha}\alpha$ in $V[G]$ and there is a well-order of $H(\alpha^+)^{V[G]}$ that is definable in $\langle H(\alpha^+)^{V[G]}, \in \rangle$ by a formula with parameters.*

Proof. It follows directly from the definition of the forcing iteration that the partial order $\vec{\mathbb{C}}_{< \alpha}$ has cardinality α . This implies $(2^{\alpha})^{V[G]} = (2^{\alpha})^V$ and we can apply Lemma 2.3 to conclude $(2^{\alpha})^{V[G][G_{\alpha}]} = (2^{\alpha})^V$. By Proposition 4.2, α is an

inaccessible cardinal in $V[\bar{G}]$ and there is an α -coding basis $\langle A, s \rangle$ in $V[\bar{G}]$ such that $\langle A, s, \mathbb{1}_{\mathbb{P}_s(A)} \rangle \in G_\alpha$. Theorem 2.9 shows that A is a Σ_1^1 -subset of ${}^\alpha\alpha$ in $V[\bar{G}][G_\alpha]$ and there is a well-order of $H(\alpha^+)^{V[\bar{G}][G_\alpha]}$ definable in $\langle H(\alpha^+)^{V[\bar{G}][G_\alpha]}, \in \rangle$ by a formula with parameters by Theorem 3.1. As above, it is easy to show that $\dot{\mathbb{C}}_{[\alpha+1, \nu]}^{\bar{G} * G_\alpha}$ adds no new α -sequences of ordinals. We can conclude $(2^\alpha)^{V[G]} = (2^\alpha)^V$, $({}^\alpha\alpha)^{V[G]} = ({}^\alpha\alpha)^{V[\bar{G} * G_\alpha]}$ and $H(\alpha^+)^{V[G]} = H(\alpha^+)^{V[\bar{G}][G_\alpha]}$. \square

Proposition 4.5. *Let κ be an infinite cardinal with the property that $\kappa \notin (\nu^+, 2^\nu]$ holds whenever ν is a singular limit of inaccessible cardinals. Given $\mu > \kappa$, $\vec{\mathbb{C}}_{<\mu}$ preserves the cardinality of κ and, if κ is regular, then $\vec{\mathbb{C}}_{<\mu}$ preserves the regularity of κ .*

Proof. By Proposition 4.3, we may assume that κ is not inaccessible. Let

$$\nu = \sup\{\alpha < \kappa \mid \alpha \text{ is an inaccessible cardinal}\}.$$

If $\nu = 0$ or ν is inaccessible, then $\nu < \kappa$, $\vec{\mathbb{C}}_{<\nu+1}$ satisfies the κ -chain condition and $\mathbb{1}_{\vec{\mathbb{C}}_{<\nu+1}} \Vdash \text{“}\dot{\mathbb{C}}_{[\nu+1, \mu]} \text{ is } <\check{\kappa}^+\text{-closed”}$ holds by Proposition 4.1.

If ν is singular and $\kappa = \nu$, then κ is a limit of inaccessible cardinals and $\vec{\mathbb{C}}_{<\mu}$ preserves the cardinality of κ by Proposition 4.3.

Let ν be singular and $\kappa = \nu^+$. Assume, toward a contradiction, that κ has cardinality less or equal to ν in some $\vec{\mathbb{C}}_{<\mu}$ -generic extension $V[G]$ of the ground model. Then there is an inaccessible cardinal α such with $\text{cof}(\kappa)^{V[G]} < \alpha < \nu$. If \bar{G} is the filter in $\vec{\mathbb{C}}_{<\alpha+1}$ induced by G , then $\text{cof}(\kappa)^{V[\bar{G}]} < \kappa$, because $\dot{\mathbb{C}}_{[\alpha+1, \mu]}^{\bar{G}}$ is $<\alpha$ -closed by Proposition 4.1. But $\vec{\mathbb{C}}_{<\alpha+1}$ satisfies the κ -chain condition, a contradiction. This shows that $\vec{\mathbb{C}}_{<\mu}$ preserves the cardinality and cofinality of ν^+ .

If ν is singular and $\kappa > 2^\nu$, then $\vec{\mathbb{C}}_{<\nu+1}$ satisfies the κ -chain condition and $\mathbb{1}_{\vec{\mathbb{C}}_{<\nu+1}} \Vdash \text{“}\dot{\mathbb{C}}_{[\nu+1, \mu]} \text{ is } <\check{\kappa}^+\text{-closed”}$ holds by Proposition 4.1. \square

5. PRESERVING SUPERCOMPACTNESS

This section is devoted to the proof of the following theorem.

Theorem 5.1. *Let γ be a cardinal with $2^\gamma = \gamma^+$ and $2^\nu \leq \gamma$, where*

$$\nu = \sup\{\alpha \leq \gamma \mid \alpha \text{ is an inaccessible cardinal}\}.$$

If κ is γ -supercompact with $\gamma = \gamma^{<\kappa}$, then

$$\mathbb{1}_{\vec{\mathbb{C}}_{<\lambda}} \Vdash \text{“}\check{\kappa} \text{ is } \check{\gamma}\text{-supercompact”}$$

holds for all $\lambda > \nu$.

Proof. By our assumptions, $\text{cof}(\gamma) \geq \kappa$ and $\nu \in [\kappa, \gamma)$ is a strong limit cardinal.

Let U be a normal ultrafilter on $\mathcal{P}_\kappa(\gamma)$. We will prove a number of claims that will allow us to show that κ is γ -supercompact in every $\vec{\mathbb{C}}_{<\nu+1}$ -generic extension of the ground model. Given $\alpha \leq \beta \in \text{On}$, we define $\vec{\mathbb{Q}}_{<\alpha} = \vec{\mathbb{C}}_{<\alpha}^{M_U}$, $\mathbb{Q}_\alpha = \dot{\mathbb{C}}_\beta^{M_U}$ and $\dot{\mathbb{Q}}_{[\alpha, \beta]} = \dot{\mathbb{C}}_{[\alpha, \beta]}^{M_U}$.

Since ν is either an inaccessible cardinal or a limit of inaccessible cardinals, we have $\vec{\mathbb{C}}_{<\alpha} \in V_\nu \subseteq M_U$ for all $\alpha < \nu$ and this shows $\vec{\mathbb{C}}_{<\nu} \in M_U$, because ${}^\gamma M_U \subseteq M_U$ holds. The definition of $\vec{\mathbb{C}}_{<\alpha}$ is absolute between V and M_U for every $\alpha \leq \nu$. Hence

elementarity implies $\vec{\mathbb{C}}_{<\nu} = \vec{\mathbb{Q}}_{<\nu}$. In particular, if \vec{G} is $\vec{\mathbb{C}}_{<\nu}$ -generic over \mathbb{V} , then \vec{G} is $\vec{\mathbb{Q}}_{<\nu}$ -generic over M_U .

Claim 1. *If \vec{G} is $\vec{\mathbb{C}}_{<\nu}$ -generic over \mathbb{V} , then $({}^\gamma M_U[\vec{G}])^{\mathbb{V}[\vec{G}]} \subseteq M_U[\vec{G}]$.*

Proof of the claim. Let $x \in \mathbb{V}[\vec{G}]$ with $x \subseteq \gamma$. We can find a $\vec{\mathbb{C}}_{<\nu}$ -nice name $\tau = \bigcup_{\alpha < \gamma} \{\check{\alpha}\} \times A_\alpha$ with $x = \tau^{\vec{G}}$. By the above remarks, we have $\vec{\mathbb{C}}_{<\nu} \subseteq {}^\nu \mathbb{V}_\nu$ and every A_α has cardinality at most $2^\nu \leq \gamma$. This shows that every A_α is an element of M_U and we also get $\langle A_\alpha \mid \alpha < \gamma \rangle \in M_U$. Hence $\tau \in M_U$ and $x = \tau^{\vec{G}} \in M_U[\vec{G}]$. We can conclude $({}^\gamma 2)^{\mathbb{V}[\vec{G}]} \subseteq M_U[\vec{G}]$.

Let $X \in \mathbb{V}[\vec{G}]$ with $X \subseteq \text{On}$ and $|X|^{\mathbb{V}[\vec{G}]} \leq \gamma$. Since $\vec{\mathbb{C}}_{<\nu}$ satisfies the γ -chain condition in \mathbb{V} , there is an $X_0 \in \mathbb{V}$ with $X \subseteq X_0$ and $|X_0|^\mathbb{V} \leq \gamma$. By our assumptions, $X_0 \in M_U$ and $|X_0|^{M_U} \leq \gamma$. Let $\langle \eta_\alpha \mid \alpha < \gamma \rangle$ be an enumeration of X_0 in M_U and $x = \{\alpha < \gamma \mid \eta_\alpha \in X\} \in \mathbb{V}[\vec{G}]$. By the above argument, $x \in M_U[\vec{G}]$ and this shows $X \in M_U[\vec{G}]$.

The argument shows $({}^\gamma \text{On})^{\mathbb{V}[\vec{G}]} \subseteq M_U[\vec{G}]$ and this implies the statement of the claim, because $M_U[\vec{G}]$ is a transitive ZFC-model with $\text{On} \subseteq M_U[\vec{G}] \subseteq \mathbb{V}[\vec{G}]$. \square

Claim 2. *If \vec{G} is $\vec{\mathbb{C}}_{<\nu}$ -generic over \mathbb{V} , then $\dot{\mathbb{C}}_\nu^{\vec{G}} = \dot{\mathbb{Q}}_\nu^{\vec{G}}$.*

Proof of the claim. If ν is not an inaccessible cardinal in \mathbb{V} , then ν is not inaccessible in M_U and both partial orders are trivial.

Now, assume that ν is inaccessible in \mathbb{V} and M_U . By Lemma 4.4, $(2^\nu)^{\mathbb{V}[\vec{G}]} = (2^\nu)^\mathbb{V} \leq \gamma$ and Claim 1 implies $\mathcal{P}({}^\nu \nu)^{\mathbb{V}[\vec{G}]} = \mathcal{P}({}^\nu \nu)^{M_U[\vec{G}]}$. This allows us to conclude $\dot{\mathbb{C}}_\nu^{\vec{G}} = \mathbb{C}_\nu^{\mathbb{V}[\vec{G}]} = \mathbb{C}_\nu^{M_U[\vec{G}]} = \dot{\mathbb{Q}}_\nu^{\vec{G}}$. \square

In particular, if G is $\vec{\mathbb{C}}_{<\nu+1}$ -generic over \mathbb{V} , then G is $\vec{\mathbb{Q}}_{<\nu+1}$ -generic over M_U .

Claim 3. *If G is $\vec{\mathbb{C}}_{<\nu+1}$ -generic over \mathbb{V} , then $({}^\gamma M_U[G])^{\mathbb{V}[G]} \subseteq M_U[G]$.*

Proof of the claim. Let \vec{G} be the filter in $\vec{\mathbb{C}}_{<\nu}$ corresponding to G and G_ν be the filter in $\dot{\mathbb{C}}_\nu^{\vec{G}}$ corresponding to G . By Proposition 3.2 and the above claims, there is a partial order \mathbb{P} in $M_U[\vec{G}]$ and $H \in M_U[G]$ such that \mathbb{P} satisfies the ν^+ -chain condition in $\mathbb{V}[\vec{G}]$, H is \mathbb{P} -generic over $\mathbb{V}[\vec{G}]$ and H induces G_ν as in (3). Every anti-chain in \mathbb{P} in $\mathbb{V}[\vec{G}]$ has cardinality at most γ in $\mathbb{V}[\vec{G}]$ and $({}^\gamma M_U[\vec{G}])^{\mathbb{V}[\vec{G}]} \subseteq M_U[\vec{G}]$, we can repeat the proof of Claim 1 and deduce the statement of the claim. \square

The proofs of the above claims show that every set of ordinals of cardinality at most γ in a $\vec{\mathbb{C}}_{<\nu+1}$ -generic extension of \mathbb{V} is covered by a set of cardinality γ in \mathbb{V} . By our assumptions, this implies that every set of ordinals of cardinality at most γ in a $\vec{\mathbb{Q}}_{<\nu+1}$ -generic extension of M_U is covered by a set of cardinality γ in M_U . In particular, forcing with $\vec{\mathbb{Q}}_{<\nu+1}$ preserves $(\gamma^+)^{M_U} = (\gamma^+)^\mathbb{V}$.

Claim 4. *If G is $\vec{\mathbb{C}}_{<\nu+1}$ -generic over \mathbb{V} , then $\dot{\mathbb{Q}}_{[\nu+1, \mu]}^G$ is $<\gamma^+$ -closed in $M_U[G]$ for all $\mu > \nu$ and the power set of $\dot{\mathbb{Q}}_{[\nu+1, j_U(\nu)]}^G$ in $M_U[G]$ has cardinality at most γ^+ in $\mathbb{V}[G]$.*

Proof of the claim. In M_U , the interval (ν, γ^+) contains no inaccessible cardinals, because ${}^\gamma M_U \subseteq M_U$ holds and no ordinal in this interval is inaccessible in \mathbb{V} . By the above remark and an application of Proposition 4.1 in M_U , we can conclude that $\dot{\mathbb{Q}}_{[\nu+1, \mu]}^G$ is $<\gamma^+$ -closed in $M_U[G]$ for all $\mu > \nu$.

By the definition of the partial order $\dot{\mathbb{C}}_{[\alpha, \beta]}$ and elementarity, the cardinality of $\dot{\mathbb{Q}}_{[\nu+1, j_U(\nu)]}^G$ in $M_U[G]$ is less or equal to the cardinality of $\vec{\mathbb{Q}}_{< j_U(\nu)}$ in M_U . The above computations and elementarity show that the cardinality of $\vec{\mathbb{Q}}_{< j_U(\nu)}$ in M_U is at most $j_U(2^\nu)$ and this ordinal is smaller or equal to $j_U(\gamma)$. If $\alpha < j_U(\gamma)$, then α is represented in M_U by a function $f : \mathcal{P}_\kappa(\gamma) \rightarrow \gamma$ contained in V . By our assumptions, $\mathcal{P}_\kappa(\gamma)$ has cardinality γ in V and there are at most 2^γ -many such functions in V . Since $2^\gamma = \gamma^+$ holds in V and $(\gamma^+)^{V[G]} = (\gamma^+)^V$, this shows that $j_U(\gamma)$ has cardinality at most γ^+ in $V[G]$. \square

Since $\vec{\mathbb{C}}_{< \nu} \in M_U$ has cardinality at most γ in V , we have $j_U \upharpoonright \vec{\mathbb{C}}_{< \nu} \in M_U$ and there is a sequence

$$\langle \dot{G}_\alpha \in (V^{\vec{\mathbb{Q}}_{< \nu}})^{M_U} \mid j_U(\kappa) \leq \alpha < j_U(\nu) \rangle$$

of names in M_U with the property that $\dot{G}_\alpha^{\vec{G}} = \{j_U(\vec{p}) \upharpoonright \alpha \mid \vec{p} \in \vec{G}\}$ for all $\alpha \in [j_U(\kappa), j_U(\nu))$ whenever \vec{G} is $\vec{\mathbb{Q}}_{< \nu}$ -generic over M_U .

Claim 5. *Let $\alpha \in [j_U(\kappa), j_U(\nu))$ be an inaccessible cardinal in M_U , H be $\vec{\mathbb{Q}}_{< \alpha}$ -generic over M_U and \vec{G} be the filter in $\vec{\mathbb{Q}}_{< \nu}$ induced by H . If $\dot{G}_\alpha^{\vec{G}} \subseteq H$ and $j_U(\vec{p})(\alpha)^H \neq \mathbb{1}_{\mathbb{C}_\alpha^{M_U[H]}}$ for some $\vec{p} \in \vec{G}$, then the following statements hold.*

- (i) *There is a unique α -coding basis $\langle A_\alpha, s_\alpha \rangle$ coding a well-order of ${}^\alpha\alpha$ in $M_U[H]$ such that for all $\vec{p} \in \vec{G}$ with $j_U(\vec{p})(\alpha)^H \neq \mathbb{1}_{\mathbb{C}_\alpha^{M_U[H]}}$ there is a $q \in \mathbb{P}_{s_\alpha}(A_\alpha)^{M_U[H]}$ with $j_U(\vec{p})(\alpha)^H = \langle A_\alpha, s_\alpha, q \rangle$.*
- (ii) *The set*

$$P_\alpha = \{q \in \mathbb{P}_{s_\alpha}(A_\alpha)^{M_U[H]} \mid (\exists \vec{p} \in \vec{G}) j_U(\vec{p})(\alpha)^H = \langle A_\alpha, s_\alpha, q \rangle\}$$

satisfies the statements (i)-(iii) of Lemma 2.10 in $M_U[H]$.

Proof of the claim. If $\vec{p} \in \vec{G}$ and $\beta < \nu$, then $\mathbb{1}_{\vec{\mathbb{C}}_{< \beta}} \Vdash \text{“}\vec{p}(\beta) \in \dot{\mathbb{C}}_\beta\text{”}$. By elementarity, we have $\mathbb{1}_{\vec{\mathbb{Q}}_{< \alpha}} \Vdash \text{“}j_U(\vec{p})(\alpha) \in \dot{\mathbb{Q}}_\alpha\text{”}$ and, by Proposition 4.2, this implies

$$Q_\alpha = \{j_U(\vec{p})(\alpha)^H \mid \vec{p} \in \vec{G}\} \subseteq \dot{\mathbb{Q}}_\alpha^H = \mathbb{C}_\alpha^{M_U[H]}.$$

Given $\vec{p}_0, \vec{p}_1 \in \vec{G}$, there is a $\vec{p} \in \vec{G}$ with $\vec{p} \leq_{\vec{\mathbb{C}}_{< \nu}} \vec{p}_0, \vec{p}_1$ and hence $\vec{p} \upharpoonright \beta \Vdash \text{“}\vec{p}(\beta) \leq_{\mathbb{P}_\beta} \vec{p}_0(\beta), \vec{p}_1(\beta)\text{”}$ for all $\beta < \nu$. Since $j_U(\vec{p}) \upharpoonright \alpha \in \dot{G}_\alpha^{\vec{G}} \subseteq H$, this argument shows that the elements of Q_α are pairwise compatible.

Pick $\vec{p}_* \in \vec{G}$ with $j_U(\vec{p}_*)(\alpha)^H \neq \mathbb{1}_{\mathbb{C}_\alpha^{M_U[H]}}$ and define $\langle A_\alpha, s_\alpha \rangle \in M_U[H]$ to be the unique α -coding basis coding a well-order of ${}^\alpha\alpha$ with $j_U(\vec{p}_*)(\alpha)^H = \langle A_\alpha, s_\alpha, q \rangle$ for some condition $q \in \mathbb{P}_{s_\alpha}(A_\alpha)^{M_U[H]}$. By the above computations, every element of Q_α is either of the form $\mathbb{1}_{\mathbb{C}_\alpha^{M_U[H]}}$ or $\langle A_\alpha, s_\alpha, q \rangle$ for some $q \in \mathbb{P}_{s_\alpha}(A_\alpha)^{M_U[H]}$.

Since \vec{G} has cardinality at most γ in $M_U[H]$, $\gamma < j_U(\kappa) \leq \alpha$ and α is regular in $M_U[H]$, we know that $\eta = \text{lub}\{\text{ht}(T_q) \mid q \in P_\alpha\} < \alpha$ and $\bigcup\{\text{dom}(f_q) \mid q \in P_\alpha\}$ has cardinality less than α in $M_U[H]$.

We show that $\eta \in \text{Lim} \cap \alpha$. Let $\vec{p} \in \vec{G}$ and $p \in \mathbb{P}_{s_\alpha}(A_\alpha)^{M_U[H]}$ with $\langle A_\alpha, s_\alpha, p \rangle = j_U(\vec{p})(\alpha)^H \neq \mathbb{1}_{\mathbb{C}_\alpha^{M_U[H]}}$. Let D be the set consisting of all conditions $\vec{q} \in \vec{\mathbb{C}}_{< \nu}$ with $\vec{q} \leq_{\vec{\mathbb{C}}_{< \nu}} \vec{p}$ and

$$\begin{aligned} \vec{q} \upharpoonright \beta \Vdash \text{“}(\forall A, s, p) [(\dot{\mathbb{C}}_\beta = \mathbb{C}_\beta \wedge \vec{p}(\beta) = \langle A, s, p \rangle \neq \mathbb{1}_{\mathbb{C}_\beta}) \\ \rightarrow (\exists \vec{p})[\vec{q}(\beta) = \langle A, s, \vec{p} \rangle \wedge \text{ht}(T_{\vec{p}}) < \text{ht}(T_{\vec{p}})]]\text{”} \end{aligned}$$

for all $\beta < \nu$. An easy inductive construction using Lemma 2.4 shows that D is dense below \vec{p} in V . If $\vec{q} \in D \cap \vec{G}$ with $j_U(\vec{q})(\alpha)^H = \langle A_\alpha, s_\alpha, q \rangle$, then $\text{ht}(T_q) > \text{ht}(T_p)$ holds in $M_U[H]$ by elementarity. This shows that η is a limit ordinal.

Finally, the conditions in P_α are pairwise compatible, because the conditions in Q_α are pairwise compatible and the first part of the claim shows that every condition in P_α belongs to a condition in Q_α . \square

In M_U , we define a sequence $\vec{q}_* = \langle \dot{q}_\alpha \in (V^{\vec{Q}_{<\alpha}})^{M_U} \mid \alpha < j_U(\nu) \rangle$ such that the following statements hold in M_U for all $\alpha < j_U(\nu)$.

- (i) If $\alpha < j_U(\kappa)$ or α is not an inaccessible cardinal, then $\mathbb{1}_{\vec{Q}_{<\alpha}} \Vdash " \dot{q}_\alpha = \dot{\mathbb{1}}_{\dot{Q}_\alpha} "$.
- (ii) If α is an inaccessible cardinal in $[j_U(\kappa), j_U(\nu))$, then \dot{q}_α is a canonical $\vec{Q}_{<\alpha}$ -name τ such that the following statements hold whenever H is $\vec{Q}_{<\alpha}$ -generic over M_U and \vec{G} is the filter in $\vec{Q}_{<\nu}$ induced by H .
 - (a) If $\dot{G}_\alpha^{\vec{G}} \subseteq H$ and $j_U(\vec{p})(\alpha)^H \neq \mathbb{1}_{\dot{Q}_\alpha^H}$ for some $\vec{p} \in \vec{G}$, then $\tau^H = \langle A_\alpha, s_\alpha, p_{P_\alpha} \rangle$, where A_α , s_α and P_α are defined as in Claim 5 and p_{P_α} is defined as in Lemma 2.10.
 - (b) Otherwise, $\tau^H = \mathbb{1}_{\dot{Q}_\alpha^H}$.

Claim 6. $\vec{q}_* \in \vec{Q}_{<j_U(\nu)}$.

Proof of the claim. Let $\alpha \in [j_U(\kappa), j_U(\nu))$ be a regular cardinal in M_U . For all $\vec{p} \in \vec{C}_{<\nu}$ there is an $\bar{\alpha}_{\vec{p}} < \alpha$ with $j_U(\vec{p})(\beta) = \dot{\mathbb{1}}_{\dot{Q}_\beta}$ for all $\bar{\alpha}_{\vec{p}} \leq \beta < \alpha$. Since $j_U''\vec{C}_{<\nu}$ is an element of M_U and has cardinality less than α in M_U , we can find an $\bar{\alpha} \in (j_U(\kappa), \alpha)$ with $j_U(\vec{p})(\beta) = \dot{\mathbb{1}}_{\dot{Q}_\beta}$ for all $\vec{p} \in \vec{C}_{<\nu}$ and $\bar{\alpha} \leq \beta < \alpha$. If $\beta \in (\bar{\alpha}, \alpha)$ is an inaccessible cardinal, H is $\vec{Q}_{<\alpha}$ -generic over M_U and \vec{G} is the filter in $\vec{Q}_{<\nu}$ induced by H , then $j_U(\vec{p})(\beta)^H = \mathbb{1}_{\dot{Q}_\beta^H}$ for all $\vec{p} \in \vec{G}$ and $\dot{q}_\beta^H = p_{P_\beta} = \mathbb{1}_{\dot{Q}_\beta^H}$ by the uniqueness of p_{P_β} . By the definition of \dot{q}_β , this shows $\dot{q}_\beta = \dot{\mathbb{1}}_{\dot{Q}_\beta}$. Therefore \vec{q}_* is a sequence with Easton support. \square

Claim 7. If H is $\vec{Q}_{<j_U(\nu)}$ -generic over M_U with $\vec{q}_* \in H$ and \vec{G} is the corresponding filter in $\vec{Q}_{<\nu}$, then $j_U''\vec{G} \subseteq H$.

Proof of the claim. Let $\alpha \in [\nu, j_U(\nu))$ and F be $\vec{Q}_{<\alpha}$ -generic over M_U with $\vec{q}_* \restriction \alpha \in F$. Assume that F induces \vec{G} in $\vec{Q}_{<\nu}$ and

$$(4) \quad \vec{q}_* \restriction [\nu, \alpha] \leq_{\dot{Q}_{[\nu, \alpha]}^{\vec{G}}} j_U(\vec{p}) \restriction [\nu, \alpha]$$

holds for all $\vec{p} \in \vec{G}$. Pick $\vec{p} \in \vec{G}$. There is a $\bar{\kappa} < \kappa$ such that $\vec{p}(\beta) = \dot{\mathbb{1}}_{\dot{Q}_\beta}$ for all $\beta \in [\bar{\kappa}, \kappa)$ and

$$j_U(\vec{p})(\beta) = \begin{cases} \vec{p}(\beta), & \text{if } \beta < \bar{\kappa}, \\ \dot{\mathbb{1}}_{\dot{Q}_\beta}, & \text{if } \bar{\kappa} \leq \beta < \nu. \end{cases}$$

by the definition of $\vec{C}_{<\nu}$. In particular, $\vec{p} \leq_{\vec{Q}_{<\nu}} j_U(\vec{p}) \restriction \nu$. By our assumption, there is a $\vec{p}_* \in \vec{G}$ with $\vec{p}_* \leq_{\vec{Q}_{<\nu}} \vec{p}$ and

$$\vec{p}_* * (\vec{q}_* \restriction [\nu, \alpha]) \leq_{\vec{Q}_{<\nu} * \dot{Q}_{[\nu, \alpha]}} i_{[\nu, \alpha]}(j_U(\vec{p}) \restriction \alpha).$$

This implies $j_U(\vec{p}) \restriction \alpha \in F$ and hence $\dot{G}_\alpha^{\vec{G}} \subseteq F$.

Next, we show that (4) holds in $M_U[G]$ for all $\vec{p} \in \vec{G}$ and $\alpha \in [\nu, j_U(\nu)]$ by induction. The case “ $\alpha = \nu$ ” is trivial and the case “ $\alpha \in \text{Lim}$ ” follows directly from the induction hypothesis.

Assume $\alpha = \bar{\alpha} + 1$ with $\bar{\alpha} \geq \nu$. We may assume that $\bar{\alpha}$ is an inaccessible cardinal in M_U . It suffices to show that

$$\vec{q}_*(\bar{\alpha})^F \leq_{\dot{\mathbb{Q}}_{\bar{\alpha}}^F} j_U(\vec{p})(\bar{\alpha})^F$$

holds in $M_U[F]$ whenever $\vec{p} \in \vec{G}$ and F is $\dot{\mathbb{Q}}_{<\bar{\alpha}}$ -generic over M_U such that $\vec{q}_* \upharpoonright \bar{\alpha} \in F$ and F induces \vec{G} in $\dot{\mathbb{Q}}_{<\nu}$. We may assume that there is a $\vec{p} \in \vec{G}$ with $j_U(\vec{p})(\bar{\alpha})^F \neq \mathbb{1}_{\dot{\mathbb{Q}}_{\bar{\alpha}}^F}$. By the induction hypothesis and the above computations, we directly get $\dot{G}_{\bar{\alpha}}^{\vec{G}} \subseteq F$. The definition of $\vec{q}_*(\bar{\alpha})$ and Claim 5 imply

$$\vec{q}_*(\bar{\alpha})^F = \langle A_\alpha, s_\alpha, p_{P_\alpha} \rangle \leq_{\dot{\mathbb{Q}}_{\bar{\alpha}}^F} j_U(\vec{p})(\bar{\alpha})^F$$

for all $\vec{p} \in \vec{G}$.

This induction shows that (4) holds if $\alpha = j_U(\nu)$ and $\vec{p} \in G$. This allows us to repeat the above computation and conclude $j_U''G \subseteq H$. \square

Claim 8. $\mathbb{1}_{\vec{\mathbb{C}}_{<\nu+1}} \Vdash \text{“}\kappa \text{ is } \check{\gamma}\text{-supercompact”}$.

Proof of the claim. Let G be $\vec{\mathbb{C}}_{<\nu+1}$ -generic over V , \vec{G} be the corresponding filter in $\vec{\mathbb{C}}_{<\nu}$ and G_ν be the corresponding filter in $\dot{\mathbb{C}}_{\nu}^{\vec{G}}$. Claim 4 combined with Claim 3 shows that there is a $\vec{H} \in V[G]$ such that $\vec{q}_* \in \vec{H}$, \vec{H} is $\dot{\mathbb{Q}}_{<j_U(\nu)}$ -generic over M_U and \vec{H} induces G in $\dot{\mathbb{Q}}_{<\nu+1}$. By Claim 7, we have $j_U''\vec{G} \subseteq \vec{H}$ and we can apply [Cum10, Proposition 9.1] to define an elementary embedding $j : V[\vec{G}] \rightarrow M_U[\vec{H}]$ extending j_U in $V[G]$ that by setting $j(\tau^{\vec{G}}) = j_U(\tau)^{\vec{H}}$ for all $\tau \in V^{\vec{\mathbb{C}}_{<\nu}}$.

We show that there is a $H_* \in V[G]$ such that H_* is $\dot{\mathbb{Q}}_{j_U(\nu)}^{\vec{H}}$ -generic over M_U and $j''G_\nu \subseteq H_*$. We may assume that ν is an inaccessible cardinal. This implies $(2^\nu)^{V[G]} = (2^\nu)^V \leq \gamma$. By Proposition 3.2, there is a ν -coding basis $\langle A, s \rangle \in V[\vec{G}]$ coding a well-order of ${}^\nu\nu$ and a filter $F_\nu \in V[G]$ such that F_ν is $\mathbb{P}_s(A)^{V[\vec{G}]}$ -generic over $V[\vec{G}]$ and F_ν induces G_ν as in (3).

By Claim 3, we have $({}^\gamma\text{On})^{V[G]} \subseteq M_U[G] \subseteq M_U[\vec{H}] \subseteq V[G]$ and this implies that $({}^\gamma M_U[\vec{H}])^{V[G]} \subseteq M_U[\vec{H}]$ holds. In particular, both $\mathbb{P}_s(A)^{V[\vec{G}]}$ and $j \upharpoonright \mathbb{P}_s(A)^{V[\vec{G}]}$ are elements of $M_U[\vec{H}]$, because $\mathbb{P}_s(A)^{V[\vec{G}]}$ has cardinality at most γ in $V[\vec{G}]$. If $j(\langle A, s \rangle) = \langle \bar{A}, \bar{s} \rangle$ and $P = j''F_\nu$, then $\langle \bar{A}, \bar{s} \rangle$ is a $j_U(\nu)$ -coding basis that codes a well-order of $j_U(\nu)$ in $M_U[\vec{H}]$, $P \subseteq \mathbb{P}_{\bar{s}}(\bar{A})^{M_U[\vec{H}]}$ and $P \in M_U[\vec{H}]$, because F_ν is an element of $M_U[\vec{H}]$. As in the proof of Claim 5, the set P satisfies the statements (i)-(iii) of Lemma 2.10 in $M_U[\vec{H}]$ and we can find a condition $p_P \in \mathbb{P}_{\bar{s}}(\bar{A})^{M_U[\vec{H}]}$ as in the statement of the Lemma.

In $M_U[\vec{H}]$, $\mathbb{P}_{\bar{s}}(\bar{A})^{M_U[\vec{H}]}$ is $<\gamma^+$ -closed and has cardinality at most $j_U(\gamma)$. By the proof of Claim 4, $j_U(\gamma)$ has cardinality at most γ^+ in $V[G]$ and there is a $F_* \in V[G]$ such that $p_P \in F_*$ and F_* is $\mathbb{P}_{\bar{s}}(\bar{A})^{M_U[\vec{H}]}$ -generic over $M_U[\vec{H}]$. If $H_* \in V[G]$ is the filter in $\mathbb{C}_{j_U(\nu)}^{M_U[\vec{H}]}$ corresponding to F_* , then H_* is $\dot{\mathbb{Q}}_{j_U(\nu)}^{\vec{H}}$ -generic over $M_U[\vec{H}]$ and our construction ensures $j''G_\nu \subseteq H_*$. Another application of [Cum10, Proposition 9.1] to define an elementary embedding $j_* : V[G] \rightarrow M_U[\vec{H}][H_*]$ in $V[G]$ that extends j . Since $({}^\gamma\text{On})^{V[G]} \subseteq M_U[\vec{H}][H_*] \subseteq V[G]$, this argument shows that κ is γ -supercompact in $V[G]$. \square

Claim 9. *If $\lambda > \nu$, then $\mathbb{1}_{\vec{\mathbb{C}}_{<\lambda}} \Vdash \text{“}\kappa \text{ is } \check{\gamma}\text{-supercompact”}$.*

Proof of the claim. Let H be $\vec{\mathbb{C}}_{<\lambda}$ -generic over V and G be the corresponding filter in $\vec{\mathbb{C}}_{<\nu+1}$. There are no inaccessible cardinals in (ν, γ^+) and the above computations show that $\vec{\mathbb{C}}_{<\nu+1}$ has the property that every set of ordinals of cardinality at most γ in a $\vec{\mathbb{C}}_{<\nu+1}$ -generic extension of the ground model is covered by a set of cardinality γ in V . By Proposition 4.1, $\dot{\mathbb{C}}_{[\nu+1, \lambda]}^G$ is $<\gamma^+$ -closed in $V[G]$.

By Claim 8, there is a normal filter U^* on $\mathcal{P}_\kappa(\gamma)$ in $V[G]$ and U^* is also a normal filter on $\mathcal{P}_\kappa(\gamma)$ in $V[H]$, because $V[H]$ is a $\dot{\mathbb{C}}_{[\nu+1, \lambda]}^G$ -generic extension of $V[G]$ and $<\gamma^+$ -closed forcing preserve normal filters on $\mathcal{P}_\kappa(\gamma)$. \square

This completes the proof of the theorem. \square

The following result due to Robert Solovay shows that, given a supercompact cardinal κ , there is a proper class of cardinals γ satisfying the assumptions of Theorem 5.1 with respect to κ . Remember that an uncountable cardinal is *strongly compact* if for any set S , every κ -complete filter on S can be extended to a κ -complete ultrafilter on S . Every supercompact cardinal is strongly compact (see [Kan03, Corollary 22.18]).

Theorem 5.2 ([Sol74, Theorem 1]). *If κ is a strongly compact cardinal and γ is a singular strong limit cardinal greater than κ , then $2^\gamma = \gamma^+$.*

Let κ be a cardinal and $\gamma_0 \geq \kappa$. There is a singular strong limit cardinal $\gamma > \gamma_0$ such that $\text{cof}(\gamma) \geq \kappa$ and there are no inaccessible cardinals in $(\gamma_0, \gamma]$. If κ is supercompact, then $2^\gamma = \gamma^+$ by Theorem 5.2 and γ satisfies the assumptions of Theorem 5.1. This proves the following statement.

Corollary 5.3. *If κ is supercompact and $\gamma \in \text{On}$, then there is a $\nu \in \text{On}$ with*

$$\mathbb{1}_{\vec{\mathbb{C}}_{<\lambda}} \Vdash \text{“}\check{\kappa} \text{ is } \check{\gamma}\text{-supercompact”}$$

for all $\lambda > \nu$. \square

6. PROOFS OF THE MAIN RESULTS

Given $\alpha \leq \beta \in \text{On}$, let $\epsilon_{\alpha, \beta} : \vec{\mathbb{C}}_{<\alpha} \longrightarrow \vec{\mathbb{C}}_{<\beta}$ denote the canonical embedding of partial orders. Let D be the class of all \vec{p} such that there is a $\beta \in \text{On}$ with $\vec{p} \in \vec{\mathbb{C}}_{<\beta}$ and $\vec{p} \neq \epsilon_{\alpha, \beta}(\vec{q})$ for all $\alpha < \beta$ and $\vec{q} \in \vec{\mathbb{C}}_{<\alpha}$. Define \mathbb{P} to be the class forcing with domain D ordered by $\vec{p} \leq_{\mathbb{P}} \vec{q}$ if there are $\alpha, \beta, \gamma \in \text{On}$ with $\alpha, \beta \leq \gamma$, $\vec{p} \in \vec{\mathbb{C}}_{<\alpha}$, $\vec{q} \in \vec{\mathbb{C}}_{<\beta}$ and $\epsilon_{\alpha, \gamma}(\vec{p}) \leq_{\vec{\mathbb{C}}_{<\gamma}} \epsilon_{\beta, \gamma}(\vec{q})$. This means that \mathbb{P} is a direct limit of the directed system $\langle \langle \vec{\mathbb{C}}_{<\alpha} \mid \alpha \in \text{On} \rangle, \langle \epsilon_{\alpha, \beta} \mid \alpha \leq \beta \in \text{On} \rangle \rangle$. Since $\vec{\mathbb{C}}_{<\alpha}$ is uniformly definable in parameter α , \mathbb{P} is definable without parameters.

Proof of Theorem 1.4. First, assume that the inaccessible cardinals are bounded in On and define

$$\nu = \sup\{\alpha \in \text{On} \mid \alpha \text{ is an inaccessible cardinal}\}.$$

We have $\mathbb{1}_{\vec{\mathbb{C}}_{<\nu+1}} \Vdash \text{“}\dot{\mathbb{C}}_{[\nu+1, \lambda]} \text{ is trivial”}$ for all $\lambda > \nu$ and this shows that \mathbb{P} is forcing equivalent to $\vec{\mathbb{C}}_{<\nu+1}$. Since ν is definable without parameters and each $\vec{\mathbb{C}}_{<\alpha}$ is definable in parameter α , the partial order $\vec{\mathbb{C}}_{<\nu+1}$ is definable without parameters. Proposition 4.3, Lemma 4.4 and Corollary 5.3 show that $\vec{\mathbb{C}}_{<\nu+1}$ satisfies the statements listed in Theorem 1.4 under this assumption.

Now, assume that there are unboundedly many inaccessible cardinals in On . Let G be \mathbb{P} -generic over V .

For each $\beta \in \text{On}$, define $G_\beta = \{\epsilon_{\alpha,\beta}(\vec{p}) \mid \alpha \leq \beta, \vec{p} \in G \cap \vec{C}_{<\alpha}\}$. Then G_β is $\vec{C}_{<\beta}$ -generic over V , $V[G]$ is the union of all $V[G_\beta]$ and G_α is the filter induced by G_β in $\vec{C}_{<\alpha}$ whenever $\alpha \leq \beta \in \text{On}$.

Claim 1. *If α is an inaccessible cardinal in V and $x \in V[G]$ is a subset of α , then $x \in V[G_{\alpha+1}]$.*

Proof of the claim. There is a $\beta > \alpha$ with $x \in V[G_\beta]$. Since $\vec{C}_{<\alpha+1}$ satisfies the α^+ -chain condition in V , we can apply Proposition 4.1 to show that $\dot{C}_{[\alpha+1,\beta]}$ is $<\alpha^+$ -closed in $V[G_{\alpha+1}]$ and this implies $x \in V[G_{\alpha+1}]$. \square

Claim 2. *Let x be an element of $V[G]$. There is an inaccessible cardinal α such that $y \in V[G_{\alpha+1}]$ for all $y \in V[G]$ with $y \subseteq x$. In particular, $V[G]$ satisfies the Power Set Axiom.*

Proof of the claim. By our assumption, we can find an inaccessible cardinal α in V such that $x \in V[G_{\alpha+1}]$ and $|x|^{V[G_{\alpha+1}]} \leq \alpha$. Let $i : x \rightarrow \alpha$ be an injection in $V[G_{\alpha+1}]$. If $y \in V[G]$ is a subset of x , then there is $\beta > \alpha$ with $y \in V[G_\beta]$. By Claim 1, we have $f''y \in V[G_{\alpha+1}]$ and therefore $y \in V[G_{\alpha+1}]$. This argument shows that $\mathcal{P}(x)^{V[G_{\alpha+1}]}$ is the power set of x in $V[G]$. \square

Claim 3. *$V[G]$ is a model of ZFC.*

Proof of the claim. Let \vec{p} be a condition in \mathbb{P} , $A \in V$ and $\langle D_a \mid a \in A \rangle$ be a V -definable sequence of dense subclasses of \mathbb{P} . There is $\alpha \in \text{On}$ with $\vec{p} \in \vec{C}_{<\alpha}$. Given $a \in A$, define $d_a = \{\vec{q} \restriction \alpha \mid (\exists \beta \geq \alpha) \vec{q} \in D_a \cap \vec{C}_{<\beta}\} \in V$. Then $\langle d_a \mid a \in A \rangle \in V$ and each d_a is predense in \mathbb{P} . This shows that \mathbb{P} is *pretame* with respect to V (see [Fri00, page 33]). By [Fri00, Lemma 2.19], this implies that $V[G]$ is a model of ZFC^- . \square

Claim 4. *Let κ be a cardinal in V with the property that there is no singular limit of inaccessible cardinals ν with $\nu^+ < \kappa \leq 2^\nu$ in V . Then κ is a cardinal in $V[G]$ and, if κ is regular in V , then κ is regular in $V[G]$.*

Proof of the claim. By Proposition 4.5, κ is a cardinal in $V[G_\mu]$ for every $\mu \in \text{On}$ and, if κ is regular in V , then κ is regular in every $V[G_\mu]$. In combination with the above remarks, this directly implies the statement of the claim. \square

Claim 5. *If κ is a supercompact cardinal in V , then κ is supercompact in $V[G]$.*

Proof of the claim. Given $\gamma \in \text{On}$, Corollary 5.3 shows that there is a $\nu \in \text{On}$ such that κ is γ -supercompact in $V[G_\beta]$ for all $\beta > \nu$. By Claim 2, there is an inaccessible cardinal α such that $\mathcal{P}(\mathcal{P}_\kappa(\gamma))^{V[G]} = \mathcal{P}(\mathcal{P}_\kappa(\gamma))^{V[G_\alpha]}$ and therefore $\mathcal{P}(\mathcal{P}_\kappa(\gamma))^{V[G_\alpha]} = \mathcal{P}(\mathcal{P}_\kappa(\gamma))^{V[G_\beta]}$ for all $\beta > \nu$. We can conclude that κ is γ -supercompact in $V[G]$. \square

Claim 6. *If α is an inaccessible cardinal in V , then α is an inaccessible cardinal in $V[G]$ and $(2^\alpha)^{V[G]} = (2^\alpha)^V$.*

Proof of the claim. By Proposition 4.3, α is an inaccessible cardinal in $V[G_{\alpha+1}]$ and Lemma 4.4 shows that $(2^\alpha)^{V[G_{\alpha+1}]} = (2^\alpha)^V$ holds. The statement of the claim follows directly from Claim 1. \square

Claim 7. *Let α be an inaccessible cardinal in V . There is a well-order of $H(\alpha^+)^{V[G]}$ that is definable in $\langle H(\alpha^+)^{V[G]}, \in \rangle$ by a formula with parameters.*

Proof of the claim. By Claim 2, there is a $\nu > \alpha$ with $H(\alpha^+)^{V[G]} = H(\alpha^+)^{V[G_\nu]}$. The statements of the Claim follows directly from Lemma 4.4. \square

This completes the proof of the theorem. \square

Proof of Theorem 1.1. Let α be an inaccessible cardinal and A be a subset of ${}^\alpha\alpha$. There is a $\vec{C}_{<\alpha}$ -name \dot{p} with the property that, whenever G is $\vec{C}_{<\alpha}$ -generic over V , then there is a α -coding basis $\langle \bar{A}, \bar{s} \rangle$ coding a well-order of ${}^\alpha\alpha$ in $V[G]$ that satisfies the following statements in $V[G]$.

- (i) $\dot{p}^G = \langle \bar{A}, \bar{s}, \mathbb{1}_{\mathbb{P}_{\bar{s}}(\bar{A})^{V[G]}} \rangle \in \dot{C}_\alpha^G$.
- (ii) There is a well-order \leq of ${}^\alpha\alpha$ witnessing that $\langle \bar{A}, \bar{s} \rangle$ codes a well-order of ${}^\alpha\alpha$ such that A is an initial segment of this order of order-type $|A|$.

Pick $\vec{p} \in \vec{C}_{<\alpha+1}$ with $\vec{p}(\alpha) = \dot{p}$. Then p is a condition in \mathbb{P} .

Let G be \mathbb{P} -generic over V with $p \in G$. For each $\beta \in \text{On}$, define G_β as in the proof of Theorem 1.4 and let $\dot{p}^{G_\alpha} = \langle \bar{A}, \bar{s}, \mathbb{1}_{\mathbb{P}_{\bar{s}}(\bar{A})^{V[G_\alpha]}} \rangle \in V[G_\alpha]$. By Claim 2 in the above proof, there is a $\nu > \alpha$ with $H(\alpha^+)^{V[G]} = H(\alpha^+)^{V[G_\nu]}$. Lemma 4.4 implies that \bar{A} is a Σ_1^1 -subset of ${}^\alpha\alpha$ in $V[G_\nu]$ and therefore also in $V[G]$. Let \leq denote the well-order of $({}^\alpha\alpha)^{V[G_\alpha]}$ produced by the above construction. Then \leq is definable in $\langle H(\alpha^+)^{V[G]}, \in \rangle$ and A is either equal to the domain of \leq or to the set of all \leq -predecessors of an element of this domain. This shows that A is definable in $\langle H(\alpha^+)^{V[G]}, \in \rangle$ by a Σ_1 -formula with parameters. \square

7. OPEN PROBLEMS

We close this paper with some open problems related to the above results.

If the *Singular Cardinal Hypothesis* holds, then forcing with the class-sized partial order constructed in Theorem 1.4 does not collapse cardinals. It is not obvious if the converse of this implication also holds.

Question 7.1. *Is it consistent that the partial order constructed in the proof of Theorem 1.4 collapses cardinals?*

Given a κ -coding basis $\langle A, s \rangle$, an easy argument shows that forcing with $\mathbb{P}_s(A)$ adds a Cohen-subset of κ . Therefore, a positive answer to the above question would follow from the existence of certain *scales* (see [Jec03, Definition 24.6]). The proof of [Hon10, Observation 4.3] contains the idea behind this approach.

As mentioned in the abstract, Theorem 1.4 can be viewed as a *boldface* version of Theorem 1.3 in the absence of the GCH. We may therefore ask whether a *lightface* version of Theorem 1.4 is possible.

Question 7.2. *Let κ be a regular uncountable cardinal κ with $\kappa = \kappa^{<\kappa}$ and $2^\kappa > \kappa^+$. Is there a cardinal preserving partial order \mathbb{P} with the property that, whenever $V[G]$ is a \mathbb{P} -generic extension of the ground model, then there is a well-order of $H(\kappa^+)^{V[G]}$ that is definable in $\langle H(\kappa^+)^{V[G]}, \in \rangle$ by a formula without parameters?*

In [Fhb], Radek Honzik and the first author use a κ^{++} -strong cardinal to produce a model with a measurable κ with $2^\kappa = \kappa^{++}$ and the property that there is a well-order of $H(\kappa^+)$ that is definable in $\langle H(\kappa^+), \in \rangle$ by a formula without parameters. It is natural to ask whether this statement is optimal.

Question 7.3. *Is it consistent that there is a measurable cardinal κ such that $2^\kappa > \kappa^{++}$ and there is a well-order of $H(\kappa^+)$ that is definable in $\langle H(\kappa^+), \in \rangle$ by a formula without parameters?*

The result mentioned above is used in [Fhb] to establish the consistency of a *definable failure of the Singular Cardinal Hypothesis*, i.e. if the existence of a κ^{++} -strong cardinal is consistent, then it is consistent that \aleph_ω is a strong limit cardinal, $2^{\aleph_\omega} = \aleph_{\omega+2}$ and there is a well-order of $H(\aleph_{\omega+1})$ that is definable in $\langle H(\aleph_{\omega+1})^{V[G]}, \in \rangle$ by a formula without parameters.

Starting from a supercompact cardinal, we can apply the *Laver preparation* (see [Lav78]) and Theorem 1.4 to produce a positive answer to the *boldface* version of Question 7.3. We may therefore ask whether the existence of stronger definable failure of the *Singular Cardinal Hypothesis* is consistent.

Question 7.4. *Is it consistent that there is a singular strong limit cardinal ν such that $2^\nu > \nu^{++}$ and there is a well-order of $H(\nu^+)$ that is definable in $\langle H(\nu^+), \in \rangle$ by a formula with parameters?*

Finally, we ask whether the existence of a definable well-order of $H(\aleph_{\omega+1})$ can be forced without applying some variation of Prikry-Forcing.

Question 7.5. *Is there a partial order \mathbb{P} with cardinality less than the least inaccessible cardinal and the property that, whenever $V[G]$ is a \mathbb{P} -generic extension of the ground model, then there is a well-order of $H(\aleph_{\omega+1})^{V[G]}$ that is definable in $\langle H(\aleph_{\omega+1})^{V[G]}, \in \rangle$ by a formula with parameters?*

REFERENCES

- [AF] David Asperó and Sy-David Friedman. Definable well-orders of H_{ω_2} and GCH. Submitted. Available at <http://www.logic.univie.ac.at/~sdf/papers/joint.aspero.omega-2.pdf>.
- [AF09] David Asperó and Sy-David Friedman. Large cardinals and locally defined well-orders of the universe. *Ann. Pure Appl. Logic*, 157(1):1–15, 2009.
- [Bau83] James E. Baumgartner. Iterated forcing. In *Surveys in set theory*, volume 87 of *London Math. Soc. Lecture Note Ser.*, pages 1–59. Cambridge Univ. Press, Cambridge, 1983.
- [Cum10] James Cummings. Iterated forcing and elementary embeddings. In *The Handbook of Set Theory (M. Foreman, A. Kanamori, and M. Magidor, eds.)*, volume 2, pages 775–884. Springer, Berlin, 2010.
- [FHa] Sy-David Friedman and Peter Holy. Condensation and large cardinals. *Fundamenta Mathematicae*, to appear. Available at <http://www.logic.univie.ac.at/~sdf/papers/joint.peter.pdf>.
- [Fhb] Sy-David Friedman and Radek Honzik. A definable failure of the singular cardinal hypothesis. *Israel Journal of Mathematics*, to appear. Available at <http://www.logic.univie.ac.at/~sdf/papers/joint.radek.def.sch.pdf>.
- [FHK] Sy-David Friedman, Tapani Hyttinen, and Vadim Kulikov. Generalised descriptive set theory and classification theory. Submitted. Available at <http://www.logic.univie.ac.at/~sdf/papers/joint.tapani.vadim.pdf>.
- [Fri00] Sy-David Friedman. *Fine structure and class forcing*, volume 3 of *de Gruyter Series in Logic and its Applications*. Walter de Gruyter & Co., Berlin, 2000.
- [Fri10] Sy-David Friedman. Forcing, combinatorics and definability. In *Proceedings of the 2009 RIMS Workshop on Combinatorial Set Theory and Forcing Theory in Kyoto, Japan, RIMS Kokyuroku No. 1686*, pages 24–40. 2010.
- [Hon10] Radek Honzik. Global singularization and the failure of SCH. *Ann. Pure Appl. Logic*, 161(7):895–915, 2010.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.

- [Kan03] Akihiro Kanamori. *The higher infinite*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2003. Large cardinals in set theory from their beginnings.
- [Lav78] Richard Laver. Making the supercompactness of κ indestructible under κ -directed closed forcing. *Israel J. Math.*, 29(4):385–388, 1978.
- [Lüc] Philipp Lücke. Σ_1^1 -definability at uncountable regular cardinals. Submitted.
- [MV93] Alan Mekler and Jouko Väänänen. Trees and Π_1^1 -subsets of ${}^{\omega_1}\omega_1$. *J. Symbolic Logic*, 58(3):1052–1070, 1993.
- [Sol74] Robert M. Solovay. Strongly compact cardinals and the GCH. In *Proceedings of the Tarski Symposium (Proc. Sympos. Pure Math., Vol. XXV, Univ. California, Berkeley, Calif., 1971)*, pages 365–372, Providence, R.I., 1974. Amer. Math. Soc.
- [TV99] Stevo Todorčević and Jouko Väänänen. Trees and Ehrenfeucht-Fraïssé games. *Ann. Pure Appl. Logic*, 100(1-3):69–97, 1999.
- [Vää95] Jouko Väänänen. Games and trees in infinitary logic: A survey. In *Quantifiers (M. Krynicki, M. Mostowski and L. Szczerba, eds.)*, pages 105–138. Kluwer Academic Publishers, 1995.
- [Vää11] Jouko Väänänen. *Models and games*, volume 132 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2011.

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, UNIVERSITÄT WIEN, WÄHRINGER STRASSE 25, 1090 WIEN, AUSTRIA

E-mail address: sdf@logic.univie.ac.at

INSTITUT FÜR MATHEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG, FACHBEREICH MATHEMATIK UND INFORMATIK, UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY

E-mail address: philipp.luecke@uni-muenster.de