

# THIN STATIONARY SETS AND DISJOINT CLUB SEQUENCES

SY-DAVID FRIEDMAN AND JOHN KRUEGER

ABSTRACT. We describe two opposing combinatorial properties related to adding clubs to  $\omega_2$ : the existence of a thin stationary subset of  $P_{\omega_1}(\omega_2)$  and the existence of a disjoint club sequence on  $\omega_2$ . A special Aronszajn tree on  $\omega_2$  implies there exists a thin stationary set. If there exists a disjoint club sequence, then there is no thin stationary set, and moreover there is a fat stationary subset of  $\omega_2$  which cannot acquire a club subset by any forcing poset which preserves  $\omega_1$  and  $\omega_2$ . We prove that the existence of a disjoint club sequence follows from Martin's Maximum and is equiconsistent with a Mahlo cardinal.

Suppose that  $S$  is a fat stationary subset of  $\omega_2$ , that is, for every club set  $C \subseteq \omega_2$ ,  $S \cap C$  contains a closed subset with order type  $\omega_1 + 1$ . A number of forcing posets have been defined which add a club subset to  $S$  and preserve cardinals under various assumptions. Abraham and Shelah [1] proved that, assuming CH, the poset consisting of closed bounded subsets of  $S$  ordered by end-extension adds a club subset to  $S$  and is  $\omega_1$ -distributive. S. Friedman [5] discovered a different poset for adding a club subset to a fat set  $S \subseteq \omega_2$  with finite conditions<sup>1</sup>. This finite club poset preserves all cardinals provided that there exists a *thin stationary subset of  $P_{\omega_1}(\omega_2)$* , that is, a stationary set  $T \subseteq P_{\omega_1}(\omega_2)$  such that for all  $\beta < \omega_2$ ,  $|\{a \cap \beta : a \in T\}| \leq \omega_1$ . This notion of stationarity appears in [9] and was discovered independently by Friedman. The question remained whether it is always possible to add a club subset to a given fat set and preserve cardinals, without any assumptions.

J. Krueger introduced a combinatorial principle on  $\omega_2$  which asserts the existence of a *disjoint club sequence*, which is a pairwise disjoint sequence  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  indexed by a stationary subset of  $\omega_2 \cap \text{cof}(\omega_1)$ , where each  $\mathcal{C}_\alpha$  is club in  $P_{\omega_1}(\alpha)$ . Krueger proved that the existence of such a sequence implies there is a fat stationary set  $S \subseteq \omega_2$  which cannot acquire a club subset by any forcing poset which preserves  $\omega_1$  and  $\omega_2$ .

We prove that a special Aronszajn tree on  $\omega_2$  implies there exists a thin stationary subset of  $P_{\omega_1}(\omega_2)$ . On the other hand assuming Martin's Maximum there exists a disjoint club sequence on  $\omega_2$ . Moreover, we have the following equiconsistency result.

**Theorem 0.1.** *Each of the following statements is equiconsistent with a Mahlo cardinal: (1) There does not exist a thin stationary subset of  $P_{\omega_1}(\omega_2)$ . (2) There exists a disjoint club sequence on  $\omega_2$ . (3) There exists a fat stationary set  $S \subseteq \omega_2$  such that any forcing poset which preserves  $\omega_1$  and  $\omega_2$  does not add a club subset to  $S$ .*

---

Date: June 2005.

2000 *Mathematics Subject Classification.* 03E35; 03E40.

Both authors were supported by FWF project number P16790-N04.

<sup>1</sup>A similar poset was defined independently by Mitchell [7]

Our proof of this theorem gives a totally different construction of the following result of Mitchell [8]: If  $\kappa$  is Mahlo in  $L$ , then there is a generic extension of  $L$  in which  $\kappa = \omega_2$  and there is no special Aronszajn tree on  $\omega_2$ . The consistency of Theorem 0.1(3) provides a negative solution to the following problem of Abraham and Shelah [1]: if  $S \subseteq \omega_2$  is fat, does there exist an  $\omega_1$ -distributive forcing poset which adds a club subset to  $S$ ?

Section 1 outlines notation and background material. In Section 2 we discuss thin stationarity and prove that a special Aronszajn tree implies the existence of a thin stationary set. In Section 3 we introduce disjoint club sequences and prove that the existence of such a sequence implies there is a fat stationary set in  $\omega_2$  which cannot acquire a club subset by any forcing poset which preserves  $\omega_1$  and  $\omega_2$ . In Section 4 we prove that Martin's Maximum implies there exists a disjoint club sequence. In Section 5 we construct a model in which there is a disjoint club sequence using an RCS iteration up to a Mahlo cardinal.

Sections 3 and 4 are due for the most part to J. Krueger. We would like to thank Boban Veličković and Mirna Džamonja for pointing out Theorem 2.3 to the authors.

## 1. PRELIMINARIES

For a set  $X$  which contains  $\omega_1$ ,  $P_{\omega_1}(X)$  denotes the collection of countable subsets of  $X$ . A set  $C \subseteq P_{\omega_1}(X)$  is *club* if it is closed under unions of countable increasing sequences and is cofinal. A set  $S \subseteq P_{\omega_1}(X)$  is *stationary* if it meets every club. If  $C \subseteq P_{\omega_1}(X)$  is club then there exists a function  $F : X^{<\omega} \rightarrow X$  such that every  $a$  in  $P_{\omega_1}(X)$  closed under  $F$  is in  $C$ . If  $F : X^{<\omega} \rightarrow P_{\omega_1}(X)$  is a function and  $Y \subseteq X$ , we say that  $Y$  is *closed under  $F$*  if for all  $\vec{\gamma}$  from  $Y^{<\omega}$ ,  $F(\vec{\gamma}) \subseteq Y$ . A partial function  $H : P_{\omega_1}(X) \rightarrow X$  is *regressive* if for all  $a$  in the domain of  $H$ ,  $H(a)$  is a member of  $a$ . *Fodor's Lemma* asserts that whenever  $S \subseteq P_{\omega_1}(X)$  is stationary and  $H : S \rightarrow X$  is a total regressive function, there is a stationary set  $S^* \subseteq S$  and a set  $x$  in  $X$  such that for all  $a$  in  $S^*$ ,  $H(a) = x$ .

If  $\kappa$  is a regular cardinal let  $\text{cof}(\kappa)$  (respectively  $\text{cof}(< \kappa)$ ) denote the class of ordinals with cofinality  $\kappa$  (respectively cofinality less than  $\kappa$ ). If  $A$  is a cofinal subset of a cardinal  $\lambda$  and  $\kappa < \lambda$ , we write for example  $A \cup \text{cof}(\kappa)$  to abbreviate  $A \cup (\lambda \cap \text{cof}(\kappa))$ .

A stationary set  $S \subseteq \kappa$  is *fat* if for every club  $C \subseteq \kappa$ ,  $S \cap C$  contains closed subsets with arbitrarily large order types less than  $\kappa$ . If  $\kappa$  is the successor of a regular uncountable cardinal  $\mu$ , this is equivalent to the statement that for every club  $C \subseteq \kappa$ ,  $S \cap C$  contains a closed subset with order type  $\mu + 1$ . In particular, if  $A \subseteq \kappa^+ \cap \text{cof}(\mu)$  is stationary then  $A \cup \text{cof}(< \mu)$  is fat.

We write  $\theta \gg \kappa$  to indicate  $\theta$  is larger than  $2^{2^{|\text{cof}(\kappa)|}}$ .

A tree  $\mathcal{T}$  is a *special Aronszajn tree on  $\omega_2$*  if:

- (1)  $\mathcal{T}$  has height  $\omega_2$  and each level has size less than  $\omega_2$ ,
- (2) each node in  $\mathcal{T}$  is an injective function  $f : \alpha \rightarrow \omega_1$  for some  $\alpha < \omega_2$ ,
- (3) the ordering on  $\mathcal{T}$  is by extension of functions, and if  $f$  is in  $\mathcal{T}$  then  $f \upharpoonright \beta$  is in  $\mathcal{T}$  for all  $\beta < \text{dom}(f)$ .

By [8] if there does not exist a special Aronszajn tree on  $\omega_2$ , then  $\omega_2$  is a Mahlo cardinal in  $L$ .

If  $V$  is a transitive model of ZFC, we say that  $W$  is an *outer model of  $V$*  if  $W$  is a transitive model of ZFC such that  $V \subseteq W$  and  $W$  has the same ordinals as  $V$ .

A forcing poset  $\mathbb{P}$  is  $\kappa$ -*distributive* if forcing with  $\mathbb{P}$  does not add any new sets of ordinals with size  $\kappa$ .

If  $\mathbb{P}$  is a forcing poset,  $\dot{a}$  is a  $\mathbb{P}$ -name, and  $G$  is a generic filter for  $\mathbb{P}$ , we write  $a$  for the set  $\dot{a}^G$ .

Martin's Maximum is the statement that whenever  $\mathbb{P}$  is a forcing poset which preserves stationary subsets of  $\omega_1$ , then for any collection  $\mathcal{D}$  of dense subsets of  $\mathbb{P}$  with  $|\mathcal{D}| \leq \omega_1$ , there is a filter  $G \subseteq \mathbb{P}$  which intersects each dense set in  $\mathcal{D}$ .

A forcing poset  $\mathbb{P}$  is *proper* if for all sufficiently large regular cardinals  $\theta > 2^{|\mathbb{P}|}$ , there is a club of countable elementary substructures  $N$  of  $\langle H(\theta), \in \rangle$  such that for all  $p$  in  $N \cap \mathbb{P}$ , there is  $q \leq p$  which is *generic for  $N$* , i.e.  $q$  forces  $N[\dot{G}] \cap \mathbf{On} = N \cap \mathbf{On}$ . If  $\mathbb{P}$  is proper then  $\mathbb{P}$  preserves  $\omega_1$  and preserves stationary subsets of  $P_{\omega_1}(\lambda)$  for all  $\lambda \geq \omega_1$ . A forcing poset  $\mathbb{P}$  is *semiproper* if the same statement holds as above except the requirement that  $q$  is generic is replaced by  $q$  being *semigeneric*, i.e.  $q$  forces  $N[\dot{G}] \cap \omega_1 = N \cap \omega_1$ . If  $\mathbb{P}$  is semigeneric then  $\mathbb{P}$  preserves  $\omega_1$  and preserves stationary subsets of  $\omega_1$ .

If  $\mathbb{P}$  is  $\omega_1$ -c.c. and  $N$  is a countable elementary substructure of  $H(\theta)$ , then  $\mathbb{P}$  forces  $N[\dot{G}] \cap \mathbf{On} = N \cap \mathbf{On}$ ; so every condition in  $\mathbb{P}$  is generic for  $N$ .

We let  ${}^{<\omega}\mathbf{On}$  denote the class of finite strictly increasing sequences of ordinals. If  $\eta$  and  $\nu$  are in  ${}^{<\omega}\mathbf{On}$ , write  $\eta \leq \nu$  if  $\eta$  is an initial segment of  $\nu$ , and write  $\eta \triangleleft \nu$  if  $\eta \leq \nu$  and  $\eta \neq \nu$ . Let  $l(\eta)$  denote the length of  $\eta$ . A set  $T \subseteq {}^{<\omega}\mathbf{On}$  is a *tree* if for all  $\eta$  in  $T$  and  $k < l(\eta)$ ,  $\eta \upharpoonright k$  is in  $T$ . A *cofinal branch of  $T$*  is a function  $b : \omega \rightarrow \kappa$  such that for all  $n < \omega$ ,  $b \upharpoonright n$  is in  $T$ .

Suppose  $I$  is an ideal on a set  $X$ . Then  $I^+$  is the collection of subsets of  $X$  which are not in  $I$ . If  $S$  is in  $I^+$  let  $I \upharpoonright S$  denote the ideal  $I \cap \mathcal{P}(S)$ . For example if  $I = NS_\kappa$ , the ideal of non-stationary subsets of  $\kappa$ , a set  $S$  is in  $I^+$  iff  $S$  is stationary. In this case  $NS_\kappa \upharpoonright S$  is the ideal of non-stationary subsets of  $S$  and  $(NS_\kappa \upharpoonright S)^+$  is the collection of stationary subsets of  $S$ .

If  $\kappa$  is regular and  $\lambda \geq \kappa$  is a cardinal, then  $\text{COLL}(\kappa, \lambda)$  is a forcing poset for collapsing  $\lambda$  to have cardinality  $\kappa$ : conditions are partial functions  $p : \kappa \rightarrow \lambda$  with size less than  $\kappa$ , ordered by extension of functions.

## 2. THIN STATIONARY SETS

Let  $T$  be a cofinal subset of  $P_{\omega_1}(\omega_2)$ . We say that  $T$  is *thin* if for all  $\beta < \omega_2$  the set  $\{a \cap \beta : a \in T\}$  has size less than  $\omega_2$ . Note that if CH holds then  $P_{\omega_1}(\omega_2)$  itself is thin. A set  $S \subseteq P_{\omega_1}(\omega_2)$  is *closed under initial segments* if for all  $a$  in  $S$  and  $\beta < \omega_2$ ,  $a \cap \beta$  is in  $S$ .

**Lemma 2.1.** *If  $S \subseteq P_{\omega_1}(\omega_2)$  is stationary and closed under initial segments, then for all uncountable  $\beta < \omega_2$ , the set  $S \cap P_{\omega_1}(\beta)$  is stationary in  $P_{\omega_1}(\beta)$ .*

*Proof.* Consider  $\beta < \omega_2$  and let  $C \subseteq P_{\omega_1}(\beta)$  be a club set. Then the set  $D = \{a \in P_{\omega_1}(\omega_2) : a \cap \beta \in C\}$  is a club subset of  $P_{\omega_1}(\omega_2)$ . Fix  $a$  in  $S \cap D$ . Since  $S$  is closed under initial segments,  $a \cap \beta$  is in  $S \cap C$ .  $\square$

**Lemma 2.2.** *If there exists a thin stationary subset of  $P_{\omega_1}(\omega_2)$ , then there is a thin stationary set  $S$  such that for all uncountable  $\beta < \omega_2$ ,  $S \cap P_{\omega_1}(\beta)$  is stationary in  $P_{\omega_1}(\beta)$ .*

*Proof.* Let  $T$  be a thin stationary set. Define  $S = \{a \cap \beta : a \in T, \beta < \omega_2\}$ . Then  $S$  is thin stationary and closed under initial segments.  $\square$

A set  $S \subseteq P_{\omega_1}(\omega_2)$  is a *local club* if there is a club set  $C \subseteq \omega_2$  such that for all uncountable  $\alpha$  in  $C$ ,  $S \cap P_{\omega_1}(\alpha)$  contains a club in  $P_{\omega_1}(\alpha)$  (see [3]). Note that local clubs are stationary.

**Theorem 2.3.** *If there is a special Aronszajn tree on  $\omega_2$ , then there is a thin local club subset of  $P_{\omega_1}(\omega_2)$ .*

*Proof.* Let  $\mathcal{T}$  be a special Aronszajn tree on  $\omega_2$ . For each  $f$  in  $\mathcal{T}$  with  $\text{dom}(f) \geq \omega_1$ , define  $S_f = \{f^{-1} \restriction i : i < \omega_1\}$ . Note that  $S_f$  is a club subset of  $P_{\omega_1}(\text{dom } f)$ . For each uncountable  $\beta < \omega_2$  define  $S_\beta = \bigcup \{S_f : f \in \mathcal{T}, \text{dom}(f) = \beta\}$ . Then  $S_\beta$  has size  $\omega_1$ . Now define  $S = \bigcup \{S_\beta : \omega_1 \leq \beta < \omega_2\}$ . Clearly  $S$  is a local club. To show  $S$  is thin, it suffices to prove that whenever  $\beta < \gamma$  are uncountable and  $a$  is in  $S_\gamma$ , then  $a \cap \beta$  is in  $S_\beta$ . Fix  $f$  in  $\mathcal{T}$  and  $i < \omega_1$  such that  $a = f^{-1} \restriction i$ . Then  $f \restriction \beta$  is in  $\mathcal{T}$ , so  $(f \restriction \beta)^{-1} \restriction i$  is in  $S_\beta$ . But  $(f \restriction \beta)^{-1} \restriction i = (f^{-1} \restriction i) \cap \beta = a \cap \beta$ .  $\square$

In later sections of the paper we will construct models in which there does not exist a thin stationary subset of  $P_{\omega_1}(\omega_2)$ . Theorem 2.3 shows that in such a model there cannot exist a special Aronszajn tree on  $\omega_2$ , so by [8]  $\omega_2$  is Mahlo in  $L$ . Mitchell [8] constructed a model in which there is no special Aronszajn tree on  $\omega_2$  by collapsing a Mahlo cardinal in  $L$  to become  $\omega_2$  with a proper forcing poset. However, in Mitchell's model the set  $(P_{\omega_1}(\kappa))^L$  is a thin stationary subset of  $P_{\omega_1}(\omega_2)$ .

**Lemma 2.4.** *Suppose  $S \subseteq P_{\omega_1}(\omega_2)$  is a local club. Then  $S$  is a local club in any outer model  $W$  with the same  $\omega_1$  and  $\omega_2$ .*

*Proof.* Let  $C$  be a club subset of  $\omega_2$  such that for every uncountable  $\alpha$  in  $C$ ,  $S \cap P_{\omega_1}(\alpha)$  contains a club in  $P_{\omega_1}(\alpha)$ . Then  $C$  remains club in  $W$ . For each uncountable  $\alpha$  in  $C$ , fix a bijection  $g_\alpha : \omega_1 \rightarrow \alpha$ . Then  $\{g_\alpha \restriction i : i < \omega_1\}$  is a club subset of  $P_{\omega_1}(\alpha)$ . By intersecting this club with  $S$ , we get a club subset of  $S \cap P_{\omega_1}(\alpha)$  of the form  $\{a_i^\alpha : i < \omega_1\}$  which is increasing and continuous. Clearly this set remains a club subset of  $P_{\omega_1}(\alpha)$  in  $W$ .  $\square$

**Proposition 2.5.** (1) *Suppose there exists a thin local club in  $P_{\omega_1}(\omega_2)$ . Then there exists a thin local club in any outer model with the same  $\omega_1$  and  $\omega_2$ .* (2) *Suppose  $\kappa$  is a cardinal such that for all  $\mu < \kappa$ ,  $\mu^\omega < \kappa$ , and assume  $\mathbb{P}$  is a proper forcing poset which collapses  $\kappa$  to become  $\omega_2$ . Then  $\mathbb{P}$  forces there is a thin stationary subset of  $P_{\omega_1}(\omega_2)$ .*

*Proof.* (1) is immediate from Lemma 2.4 and the absoluteness of thinness. (2) Let  $G$  be generic for  $\mathbb{P}$  over  $V$  and work in  $V[G]$ . Since  $\mathbb{P}$  is proper,  $\omega_1$  is preserved and the set  $S = (P_{\omega_1}(\kappa))^V$  is stationary in  $P_{\omega_1}(\omega_2)$ . We claim that  $S$  is thin. If  $\beta < \omega_2$  then  $\{a \cap \beta : a \in S\} = (P_{\omega_1}(\beta))^V$ . By the assumption on  $\kappa$ , there is  $\xi < \kappa$  and a bijection  $f : \xi \rightarrow (P_{\omega_1}(\beta))^V$  in  $V$ . In  $V[G]$  there is a surjection of  $\omega_1$  onto  $\xi$  and hence a surjection of  $\omega_1$  onto  $\{a \cap \beta : a \in S\}$ .  $\square$

As we mentioned above, if CH holds then the set  $P_{\omega_1}(\omega_2)$  itself is thin. We show on the other hand that if CH fails then no club subset of  $P_{\omega_1}(\omega_2)$  is thin. The proof is actually due to Baumgartner and Taylor [2] who proved that for any club set  $C \subseteq P_{\omega_1}(\omega_2)$ , there is a countable set  $A \subseteq \omega_2$  such that  $C \cap \mathcal{P}(A)$  has size at least  $2^\omega$ . Their method of proof, which is described in the next lemma, is key to several of our results later in the paper.

**Lemma 2.6.** *Suppose  $Z$  is a stationary subset of  $\omega_2 \cap \text{cof}(\omega)$  and for each  $\alpha$  in  $Z$ ,  $M_\alpha$  is a countable cofinal subset of  $\alpha$ . Then there is a sequence  $\langle Z_s, \xi_s : s \in {}^{<\omega}2 \rangle$  satisfying:*

- (1) each  $Z_s$  is a stationary subset of  $Z$ ,
- (2) if  $s \leq t$  then  $Z_t \subseteq Z_s$ ,
- (3) if  $\alpha$  is in  $Z_s$  then  $\xi_s$  is in  $M_\alpha$ ,
- (4) if  $\alpha$  is in  $Z_{s \smallfrown 0}$  and  $\beta$  is in  $Z_{s \smallfrown 1}$ , then  $\xi_{s \smallfrown 0}$  is not in  $M_\beta$  and  $\xi_{s \smallfrown 1}$  is not in  $M_\alpha$ .

*Proof.* Let  $Z_{\langle \rangle} = Z$  and  $\xi_{\langle \rangle}$  is undefined. Suppose  $Z_s$  is given. Define  $X_s$  as the set of  $\xi$  in  $\omega_2$  such that the set  $\{\alpha \in Z_s : \xi \in M_\alpha\}$  is stationary. A straightforward argument using Fodor's Lemma shows that  $X_s$  is unbounded in  $\omega_2$ . For each  $\alpha$  in  $Z_s$  such that  $X_s \cap \alpha$  has size  $\omega_1$ , there exists  $\xi < \alpha$  in  $X_s$  such that  $\xi$  is not in  $M_\alpha$ . By Fodor's Lemma there is a stationary set  $Z'_{s \smallfrown 1} \subseteq Z_s$  and  $\xi_{s \smallfrown 0}$  in  $X_s$  such that for all  $\alpha$  in  $Z'_{s \smallfrown 1}$ ,  $\xi_{s \smallfrown 0}$  is not in  $M_\alpha$ . Let  $Z'_{s \smallfrown 0}$  denote the set of  $\alpha$  in  $Z_s$  such that  $\xi_{s \smallfrown 0}$  is in  $M_\alpha$ , which is stationary since  $\xi_{s \smallfrown 0}$  is in  $X_s$ . Now define  $Y_s$  as the set of  $\xi$  in  $\omega_2$  such that  $\{\alpha \in Z'_{s \smallfrown 1} : \xi \in M_\alpha\}$  is stationary. Then  $Y_s$  is unbounded in  $\omega_2$ . So for each  $\alpha$  in  $Z'_{s \smallfrown 0}$  such that  $Y_s \cap \alpha$  has size  $\omega_1$ , there is  $\xi < \alpha$  in  $Y_s$  which is not in  $M_\alpha$ . By Fodor's Lemma there is  $\xi_{s \smallfrown 1}$  in  $Y_s$  and  $Z_{s \smallfrown 0} \subseteq Z'_{s \smallfrown 0}$  stationary such that for all  $\alpha$  in  $Z_{s \smallfrown 0}$ ,  $\xi_{s \smallfrown 1}$  is not in  $M_\alpha$ . Now define  $Z_{s \smallfrown 1}$  as the set of  $\alpha$  in  $Z'_{s \smallfrown 1}$  such that  $\xi_{s \smallfrown 1}$  is in  $M_\alpha$ .  $\square$

**Theorem 2.7.** *Assume CH fails. Then for any club set  $C \subseteq P_{\omega_1}(\omega_2)$ ,  $C$  is not thin.*

*Proof.* Let  $F : \omega_2^{<\omega} \rightarrow \omega_2$  be a function such that any  $a$  in  $P_{\omega_1}(\omega_2)$  closed under  $F$  is in  $C$ . Let  $Z$  be the stationary set of  $\alpha$  in  $\omega_2 \cap \text{cof}(\omega)$  closed under  $F$ . For each  $\alpha$  in  $Z$  fix a countable set  $M_\alpha \subseteq \alpha$  such that  $\sup(M_\alpha) = \alpha$  and  $M_\alpha$  is closed under  $F$ . Fix a sequence  $\langle Z_s, \xi_s : s \in {}^{<\omega}2 \rangle$  as described in Lemma 2.6.

For each function  $f : \omega \rightarrow 2$  define  $b_f = cl_F(\{\xi_{f \smallfrown n} : n < \omega\})$ . Then  $b_f$  is in  $C$ . Note that if  $n < \omega$  and  $\alpha$  is in  $Z_{f \smallfrown n}$ , then  $cl_F(\{\xi_{f \smallfrown m} : m \leq n\}) \subseteq M_\alpha$ . For by Lemma 2.6(2), for  $m \leq n$ ,  $Z_{f \smallfrown n} \subseteq Z_{f \smallfrown m}$ . So  $\alpha$  is in  $Z_{f \smallfrown m}$ , and hence  $\xi_{f \smallfrown m}$  is in  $M_\alpha$  by (3). But  $M_\alpha$  is closed under  $F$ .

Let  $\gamma = \sup(\{\xi_s + 1 : s \in {}^{<\omega}2\})$ . Since  ${}^{<\omega}2$  has size  $\omega$ ,  $\gamma$  is less than  $\omega_2$ . We claim that for distinct  $f$  and  $g$ ,  $b_f \cap \gamma \neq b_g \cap \gamma$ . Let  $n < \omega$  be least such that  $f(n) \neq g(n)$ . If  $b_f \cap \gamma = b_g \cap \gamma$ , then there is  $k > n$  such that  $\xi_{g \smallfrown (n+1)}$  is in  $cl_F(\{\xi_{f \smallfrown m} : m \leq k\})$ . Fix  $\alpha$  in  $Z_{f \smallfrown k}$ . By the last paragraph,  $\xi_{g \smallfrown (n+1)}$  is in  $M_\alpha$ . But  $\alpha$  is in  $Z_{f \smallfrown (n+1)}$  by (2), which contradicts (4).  $\square$

Let  $\kappa$  be an uncountable cardinal. The *Weak Reflection Principle at  $\kappa$*  is the statement that whenever  $S$  is a stationary subset of  $P_{\omega_1}(\kappa)$ , there is a set  $Y$  in  $P_{\omega_2}(\kappa)$  such that  $\omega_1 \subseteq Y$  and  $S \cap P_{\omega_1}(Y)$  is stationary in  $P_{\omega_1}(Y)$ . Martin's Maximum implies the Weak Reflection Principle holds for all uncountable cardinals  $\kappa$  [4]. The Weak Reflection Principle at  $\omega_2$  is equivalent to the statement that for every stationary set  $S \subseteq P_{\omega_1}(\omega_2)$ , there is a stationary set of uncountable  $\beta < \omega_2$  such that  $S \cap P_{\omega_1}(\beta)$  is stationary in  $P_{\omega_1}(\beta)$ . This is equivalent to the statement that every local club subset of  $P_{\omega_1}(\omega_2)$  contains a club. The Weak Reflection Principle at  $\omega_2$  is equiconsistent with a weakly compact cardinal [3].

**Corollary 2.8.** *Suppose CH fails and there is a special Aronszajn tree on  $\omega_2$ . Then the Weak Reflection Principle at  $\omega_2$  fails.*

*Proof.* By Theorems 2.3 and 2.7, there is a thin local club subset of  $P_{\omega_1}(\omega_2)$  which is not club. Hence the Weak Reflection Principle at  $\omega_2$  fails.  $\square$

In Sections 4 and 5 we describe models in which there is no thin stationary subset of  $P_{\omega_1}(\omega_2)$ . On the other hand S. Friedman proved there always exists a thin cofinal set.

**Theorem 2.9** (Friedman). *There exists a thin cofinal subset of  $P_{\omega_1}(\omega_2)$ .*

*Proof.* We construct by induction a sequence  $\langle S_\alpha : \omega_1 \leq \alpha < \omega_2 \rangle$  satisfying the properties: (1) each  $S_\alpha$  is a cofinal subset of  $P_{\omega_1}(\alpha)$  with size  $\omega_1$ , (2) for uncountable  $\beta < \gamma$ , if  $a$  is in  $S_\gamma$  then  $a \cap \beta$  is in  $\bigcup \{S_\alpha : \omega_1 \leq \alpha \leq \beta\}$ , and (3) if  $\beta < \gamma < \omega_2$ ,  $a$  is in  $P_{\omega_1}(\gamma)$ , and  $a \cap \beta$  is in  $S_\beta$ , then there is  $b$  in  $S_\gamma$  such that  $a \subseteq b$  and  $a \cap \beta = b \cap \beta$ .

Let  $S_{\omega_1} = \omega_1$ . Given  $S_\alpha$ , let  $S_{\alpha+1}$  be the collection  $\{b \cup \{\alpha\} : b \in S_\alpha\}$ . Conditions (1), (2), and (3) follow by induction. Suppose  $\gamma < \omega_2$  is an uncountable limit ordinal and  $S_\alpha$  is defined for all uncountable  $\alpha < \gamma$ . If  $\text{cf}(\gamma) = \omega_1$  then let  $S_\gamma = \bigcup \{S_\alpha : \omega_1 \leq \alpha < \gamma\}$ . The required conditions follow by induction.

Assume  $\text{cf}(\gamma) = \omega$ . Fix an increasing sequence of uncountable ordinals  $\langle \gamma_n : n < \omega \rangle$  unbounded in  $\gamma$ . Let  $T_\gamma$  be some cofinal subset of  $P_{\omega_1}(\gamma)$  with size  $\omega_1$ . Fix  $n < \omega$ . For each  $x$  in  $T_\gamma$  and  $a$  in  $S_{\gamma_n}$  define a set  $b(a, x, n)$  in  $P_{\omega_1}(\gamma)$  inductively as follows. Let  $b(a, x, n) \cap \gamma_n = a$ . Given  $b(a, x, n) \cap \gamma_m$  in  $S_{\gamma_m}$  for some  $m \geq n$ , apply condition (3) to  $\gamma_m, \gamma_{m+1}$ , and the set

$$(b(a, x, n) \cap \gamma_m) \cup ((x \cap [\gamma_m, \gamma_{m+1}))$$

to find  $y$  in  $S_{\gamma_{m+1}}$  such that  $y \cap \gamma_m = b(a, x, n) \cap \gamma_m$  and  $x \cap [\gamma_m, \gamma_{m+1}) \subseteq y$ . Let  $b(a, x, n) \cap \gamma_{m+1} = y$ . This completes the definition of  $b(a, x, n)$ . Clearly  $b(a, x, n) \cap \gamma_n = a$ ,  $x \setminus \gamma_n \subseteq b(a, x, n)$ , and for all  $k \geq n$ ,  $b(a, x, n) \cap \gamma_k$  is in  $S_{\gamma_k}$ .

Now define  $S_\gamma = \{b(a, x, n) : n < \omega, a \in S_{\gamma_n}, x \in T_\gamma\}$ . We verify conditions (1), (2), and (3). Clearly  $S_\gamma$  has size  $\omega_1$ . Let  $\beta < \gamma$  and consider  $b(a, x, n)$  in  $S_\gamma$ . Fix  $k > n$  such that  $\beta < \gamma_k$ . Then  $b(a, x, n) \cap \gamma_k$  is in  $S_{\gamma_k}$ . So by induction  $b(a, x, n) \cap \beta$  is in  $\bigcup \{S_\alpha : \omega_1 \leq \alpha \leq \beta\}$ . Now assume  $a$  is in  $P_{\omega_1}(\gamma)$ ,  $\beta < \gamma$ , and  $a \cap \beta$  is in  $S_\beta$ . Choose  $x$  in  $T_\gamma$  such that  $a \subseteq x$ . Fix  $k$  such that  $\beta < \gamma_k$ . By the induction hypothesis there is  $a'$  in  $S_{\gamma_k}$  such that  $a \cap \gamma_k \subseteq a'$  and  $a' \cap \beta = a \cap \beta$ . Let  $c = b(a', x, k)$ . Then  $c$  is in  $S_\gamma$ ,  $c \cap \beta = (c \cap \gamma_k) \cap \beta = a' \cap \beta = a \cap \beta$ , and  $a \subseteq c$ .

To prove  $S_\gamma$  is cofinal consider  $a$  in  $P_{\omega_1}(\gamma)$ . Fix  $x$  in  $T_\gamma$  such that  $a \subseteq x$ . By induction  $S_{\gamma_0}$  is cofinal in  $P_{\omega_1}(\gamma_0)$ . So let  $y$  be in  $S_{\gamma_0}$  such that  $x \cap \gamma_0 \subseteq y$ . Then  $a$  is a subset of  $b(y, x, 0)$ .

Now define  $S = \bigcup \{S_\beta : \omega_1 \leq \beta < \omega_2\}$ . Conditions (1) and (2) imply that  $S$  is thin and cofinal in  $P_{\omega_1}(\omega_2)$ .  $\square$

### 3. DISJOINT CLUB SEQUENCES

We introduce a combinatorial property of  $\omega_2$  which implies there does not exist a thin stationary subset of  $P_{\omega_1}(\omega_2)$ . This property follows from Martin's Maximum and is equiconsistent with a Mahlo cardinal. It implies there exists a fat stationary subset of  $\omega_2$  which cannot acquire a club subset by any forcing poset which preserves  $\omega_1$  and  $\omega_2$ .

**Definition 3.1.** *A disjoint club sequence on  $\omega_2$  is a sequence  $\langle C_\alpha : \alpha \in A \rangle$  such that  $A$  is a stationary subset of  $\omega_2 \cap \text{cof}(\omega_1)$ , each  $C_\alpha$  is a club subset of  $P_{\omega_1}(\alpha)$ , and  $C_\alpha \cap C_\beta$  is empty for all  $\alpha < \beta$  in  $A$ .*

**Proposition 3.2.** *Suppose there is a disjoint club sequence on  $\omega_2$ . Then there does not exist a thin stationary subset of  $P_{\omega_1}(\omega_2)$ .*

*Proof.* Let  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  be a disjoint club sequence. Suppose for a contradiction there exists a thin stationary set. By Lemma 2.2 fix a thin stationary set  $T \subseteq P_{\omega_1}(\omega_2)$  such that for all uncountable  $\beta < \omega_2$ ,  $T \cap P_{\omega_1}(\beta)$  is stationary in  $P_{\omega_1}(\beta)$ . Then for each  $\beta$  in  $A$  we can choose a set  $a_\beta$  in  $\mathcal{C}_\beta \cap T$ . Since  $\text{cf}(\beta) = \omega_1$ ,  $\sup(a_\beta) < \beta$ . By Fodor's Lemma there is a stationary set  $B \subseteq A$  and a fixed  $\gamma < \omega_2$  such that for all  $\beta$  in  $B$ ,  $\sup(a_\beta) = \gamma$ . If  $\alpha < \beta$  are in  $B$ , then  $a_\alpha \neq a_\beta$  since  $\mathcal{C}_\alpha \cap \mathcal{C}_\beta$  is empty. So the set  $\{a_\beta : \beta \in B\}$  witnesses that  $T$  is not thin, which is a contradiction.  $\square$

**Lemma 3.3.** *Suppose there is a disjoint club sequence  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  on  $\omega_2$ . Let  $W$  be an outer model with the same  $\omega_1$  and  $\omega_2$  in which  $A$  is still stationary. Then there is a disjoint club sequence  $\langle \mathcal{D}_\alpha : \alpha \in A \rangle$  in  $W$ .*

*Proof.* By the proof of Lemma 2.4, each  $\mathcal{C}_\alpha$  contains a club set  $\mathcal{D}_\alpha$  in  $W$ . Since  $\omega_1$  is preserved, each  $\alpha$  in  $A$  still has cofinality  $\omega_1$ .  $\square$

**Theorem 3.4.** *Suppose  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  is a disjoint club sequence on  $\omega_2$ . Then  $A \cup \text{cof}(\omega)$  does not contain a club.*

*Proof.* Suppose for a contradiction that  $A \cup \text{cof}(\omega)$  contains a club. Without loss of generality  $2^{\omega_1} = \omega_2$ . Otherwise work in a generic extension  $W$  by  $\text{COLL}(\omega_2, 2^{\omega_1})$ : in  $W$  the set  $A \cup \text{cof}(\omega)$  contains a club and by Lemma 3.3 there is a disjoint club sequence  $\langle \mathcal{D}_\alpha : \alpha \in A \rangle$ .

Since  $2^{\omega_1} = \omega_2$ ,  $H(\omega_2)$  has size  $\omega_2$ . Fix a bijection  $h : H(\omega_2) \rightarrow \omega_2$ . Let  $\mathcal{A}$  denote the structure  $\langle H(\omega_2), \in, h \rangle$ . Define  $B$  as the set of  $\alpha$  in  $\omega_2 \cap \text{cof}(\omega_1)$  such that there exists an increasing and continuous sequence  $\langle N_i : i < \omega_1 \rangle$  of countable elementary substructures of  $\mathcal{A}$  such that:

- (1) for  $i < \omega_1$ ,  $N_i$  is in  $N_{i+1}$ ,
- (2) the set  $\{N_i \cap \omega_2 : i < \omega_1\}$  is club in  $P_{\omega_1}(\alpha)$ .

We claim that  $B$  is stationary in  $\omega_2$ . To prove this let  $C \subseteq \omega_2$  be club. Let  $\mathcal{B}$  be the expansion of  $\mathcal{A}$  by the function  $\alpha \mapsto \min(C \setminus \alpha)$ . Define by induction an increasing and continuous sequence  $\langle N_i : i < \omega_1 \rangle$  of elementary substructures of  $\mathcal{B}$  such that for all  $i < \omega_1$ ,  $N_i$  is in  $N_{i+1}$ . Let  $N = \bigcup \{N_i : i < \omega_1\}$ . Then  $\omega_1 \subseteq N$  so  $N \cap \omega_2$  is an ordinal. Write  $\alpha = N \cap \omega_2$ . Then  $\alpha$  is in  $C$  and  $\{N_i \cap \omega_2 : i < \omega_1\}$  is club in  $P_{\omega_1}(\alpha)$ . So  $\alpha$  is in  $B \cap C$ .

Since  $A \cup \text{cof}(\omega)$  contains a club,  $A \cap B$  is stationary. For each  $\alpha$  in  $A \cap B$  fix a sequence  $\langle N_i^\alpha : i < \omega_1 \rangle$  as described in the definition of  $B$ . Then  $\{N_i^\alpha \cap \omega_2 : i < \omega_1\} \cap \mathcal{C}_\alpha$  is club in  $P_{\omega_1}(\alpha)$ . So there exists a club set  $c_\alpha \subseteq \omega_1$  such that  $\{N_i^\alpha \cap \omega_2 : i \in c_\alpha\}$  is club and is a subset of  $\mathcal{C}_\alpha$ . Write  $i_\alpha = \min(c_\alpha)$  and let  $d_\alpha = c_\alpha \setminus \{i_\alpha\}$ .

Define  $S = \{N_i^\alpha : \alpha \in A \cap B, i \in d_\alpha\}$ . If  $N$  is in  $S$  then there is a unique pair  $\alpha$  in  $A \cap B$  and  $i$  in  $d_\alpha$  such that  $N = N_i^\alpha$ . For if  $N = N_i^\alpha = N_j^\beta$ , then  $N \cap \omega_2$  is in  $\mathcal{C}_\alpha \cap \mathcal{C}_\beta$ , so  $\alpha = \beta$ . Clearly then  $i = j$ . Also note that if  $N_i^\alpha$  is in  $S$  then  $N_i^\alpha$  is in  $N_i^\alpha$ . So the function  $H : S \rightarrow H(\omega_2)$  defined by  $H(N_i^\alpha) = N_{i_\alpha}^\alpha$  is well-defined and regressive.

We claim that  $S$  is stationary in  $P_{\omega_1}(H(\omega_2))$ . To prove this let  $F : H(\omega_2)^{<\omega} \rightarrow H(\omega_2)$  be a function. Define  $G : \omega_2^{<\omega} \rightarrow \omega_2$  by letting  $G(\alpha_0, \dots, \alpha_n)$  be equal to  $h(F(h^{-1}(\alpha_0), \dots, h^{-1}(\alpha_n)))$ . Let  $E$  be the club set of  $\alpha$  in  $\omega_2$  closed under  $G$ . Fix  $\alpha$  in  $E \cap A \cap B$ . Then there is  $i$  in  $d_\alpha$  such that  $N_i^\alpha \cap \omega_2$  is closed under  $G$ . We claim

that  $N_i^\alpha$  is closed under  $F$ . Given  $a_0, \dots, a_n$  in  $N_i^\alpha$ , the ordinals  $h(a_0), \dots, h(a_n)$  are in  $N_i^\alpha \cap \omega_2$ . So  $\gamma = G(h(a_0), \dots, h(a_n)) = h(F(a_0, \dots, a_n))$  is in  $N_i^\alpha \cap \omega_2$ . Therefore  $h^{-1}(\gamma) = F(a_0, \dots, a_n)$  is in  $N_i^\alpha$ .

Since  $S$  is stationary and  $H : S \rightarrow H(\omega_2)$  is regressive, there is a stationary set  $S^* \subseteq S$  and a fixed  $N$  such that for all  $N_i^\alpha$  in  $S^*$ ,  $H(N_i^\alpha) = N$ . The set  $S^*$ , being stationary, must have size  $\omega_2$ . So there are distinct  $\alpha$  and  $\beta$  such that for some  $i$  in  $d_\alpha$  and  $j$  in  $d_\beta$ ,  $N_i^\alpha$  and  $N_j^\beta$  are in  $S^*$ . Then  $N = N_{i_\alpha}^\alpha = N_{j_\beta}^\beta$ . So  $N \cap \omega_2$  is in  $\mathcal{C}_\alpha \cap \mathcal{C}_\beta$ , which is a contradiction.  $\square$

Abraham and Shelah [1] asked the following question: Assume that  $A$  is a stationary subset of  $\omega_2 \cap \text{cof}(\omega_1)$ . Does there exist an  $\omega_1$ -distributive forcing poset which adds a club subset to  $A \cup \text{cof}(\omega)$ ? We answer this question in the negative.

**Corollary 3.5.** *Assume that  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  is a disjoint club sequence. Let  $W$  be an outer model of  $V$  with the same  $\omega_1$  and  $\omega_2$ . Then in  $W$ ,  $A \cup \text{cof}(\omega)$  does not contain a club subset.*

*Proof.* If  $A$  remains stationary in  $W$ , then by Lemma 3.3 there is a disjoint club sequence  $\langle \mathcal{D}_\alpha : \alpha \in A \rangle$  in  $W$ . By Theorem 3.4  $A \cup \text{cof}(\omega)$  does not contain a club in  $W$ .  $\square$

#### 4. MARTIN'S MAXIMUM

In this section we prove that Martin's Maximum implies there exists a disjoint club sequence on  $\omega_2$ . We apply MM to the poset for adding a Cohen real and then forcing a continuous  $\omega_1$ -chain through  $P_{\omega_1}(\omega_2) \setminus V$ .

**Theorem 4.1** (Krueger). *Martin's Maximum implies there exists a disjoint club sequence on  $\omega_2$ .*

We will use the following theorem from [1].

**Theorem 4.2.** *Suppose  $\mathbb{P}$  is  $\omega_1$ -c.c. and adds a real. Then  $\mathbb{P}$  forces that  $(P_{\omega_1}(\omega_2) \setminus V)$  is stationary in  $P_{\omega_1}(\omega_2)$ .*

Note: Gitik [6] proved that the conclusion of Theorem 4.2 holds for any outer model of  $V$  which contains a new real and computes the same  $\omega_1$ .

Suppose that  $S$  is a stationary subset of  $P_{\omega_1}(\omega_2)$ . Following [3] we define a forcing poset  $\mathbb{P}(S)$  which adds a continuous  $\omega_1$ -chain through  $S$ . A condition in  $\mathbb{P}(S)$  is a countable increasing and continuous sequence  $\langle a_i : i \leq \beta \rangle$  of elements from  $S$ , where for each  $i < \beta$ ,  $a_i \cap \omega_1 < a_{i+1} \cap \omega_1$ . The ordering on  $\mathbb{P}(S)$  is by extension of sequences.

**Proposition 4.3.** *If  $S \subseteq P_{\omega_1}(\omega_2)$  is stationary then  $\mathbb{P}(S)$  is  $\omega$ -distributive.*

*Proof.* Suppose  $p$  forces  $\dot{f} : \omega \rightarrow \mathbf{On}$ . Let  $\theta \gg \omega_2$  be a regular cardinal such that  $\dot{f}$  is in  $H(\theta)$ . Since  $S$  is stationary, we can fix a countable elementary substructure  $N$  of the model

$$\langle H(\theta), \in, S, \mathbb{P}(S), p, \dot{f} \rangle$$

such that  $N \cap \omega_2$  is in  $S$ . Let  $\langle D_n : n < \omega \rangle$  be an enumeration of all the dense subsets of  $\mathbb{P}(S)$  in  $N$ . Inductively define a decreasing sequence  $\langle p_n : n < \omega \rangle$  of elements of  $N \cap \mathbb{P}$  such that  $p_0 = p$  and  $p_{n+1}$  is a refinement of  $p_n$  in  $D_n \cap N$ . Write  $\bigcup \{p_n : n < \omega\} = \langle b_i : i < \gamma \rangle$ . Clearly  $\bigcup \{b_i : i < \gamma\} = N \cap \omega_2$ . Since  $N \cap \omega_2$  is in  $S$ ,



the sequence  $\langle b_i : i < \gamma \rangle \cup \{\langle \gamma, N \cap \omega_2 \rangle\}$  is a condition below  $p$  which decides  $\dot{f}(n)$  for all  $n < \omega$ .  $\square$

**Theorem 4.4.** *Suppose  $\mathbb{P}$  is an  $\omega_1$ -c.c. forcing poset which adds a real. Let  $\dot{S}$  be a name such that  $\mathbb{P}$  forces  $\dot{S} = (P_{\omega_1}(\omega_2) \setminus V)$ . Then  $\mathbb{P} * \mathbb{P}(\dot{S})$  preserves stationary subsets of  $\omega_1$ .*

*Proof.* By Theorem 4.2 and Proposition 4.3, the poset  $\mathbb{P} * \mathbb{P}(\dot{S})$  preserves  $\omega_1$ . Let  $A$  be a stationary subset of  $\omega_1$  in  $V$ . Suppose  $p * \dot{q}$  is a condition in  $\mathbb{P} * \mathbb{P}(\dot{S})$  which forces  $\dot{C}$  is a club subset of  $\omega_1$ .

Let  $G$  be a generic filter for  $\mathbb{P}$  over  $V$  which contains  $p$ . In  $V[G]$  fix a regular cardinal  $\theta \gg \omega_2$  and let

$$\mathcal{A} = \langle H(\theta), \in, A, S, q, \dot{C} \rangle.$$

Fix a Skolem function  $F : H(\theta)^{<\omega} \rightarrow H(\theta)$  for  $\mathcal{A}$ . Define  $F^* : \omega_2^{<\omega} \rightarrow P_{\omega_1}(\omega_2)$  by letting

$$F^*(\alpha_0, \dots, \alpha_n) = cl_F(\{\alpha_0, \dots, \alpha_n\}) \cap \omega_2.$$

Since  $\mathbb{P}$  is  $\omega_1$ -c.c. there is a function  $H : \omega_2^{<\omega} \rightarrow P_{\omega_1}(\omega_2)$  in  $V$  such that for all  $\vec{\alpha}$  in  $\omega_2^{<\omega}$ ,  $F^*(\vec{\alpha}) \subseteq H(\vec{\alpha})$ . Let  $Z^*$  be the stationary set of  $\alpha$  in  $\omega_2 \cap \text{cof}(\omega)$  closed under  $H$ .

Working in  $V$ , since  $A$  is stationary we can fix for each  $\alpha$  in  $Z^*$  a countable cofinal set  $M_\alpha \subseteq \alpha$  closed under  $H$  with  $M_\alpha \cap \omega_1$  in  $A$ . By Fodor's Lemma there is  $Z \subseteq Z^*$  stationary and  $\delta$  in  $A$  such that for all  $\alpha$  in  $Z$ ,  $M_\alpha \cap \omega_1 = \delta$ . Fix a sequence  $\langle \xi_s, Z_s : s \in {}^{<\omega}2 \rangle$  satisfying conditions (1)–(4) of Lemma 2.6.

Let  $f : \omega \rightarrow 2$  be a function in  $V[G] \setminus V$ . For each  $n < \omega$  let  $M_n$  denote  $cl_H(\delta \cup \{\xi_{f \upharpoonright m} : m \leq n\})$ . Define  $M = \bigcup \{M_n : n < \omega\}$ . Note that  $M$  is closed under  $H$  and hence it is closed under  $F^*$ . Therefore  $N = cl_F(M)$  is an elementary substructure of  $\mathcal{A}$  such that  $N \cap \omega_2 = M$ .

As in the proof of Theorem 2.7, for all  $n < \omega$ , if  $\alpha$  is in  $Z_{f \upharpoonright n}$  then  $M_n \subseteq M_\alpha$ . Note that  $M \cap \omega_1 = \delta$ . For if  $\gamma$  is in  $M \cap \omega_1$ , there is  $n < \omega$  such that  $\gamma$  is in  $M_n$ . Fix  $\alpha$  in  $Z_{f \upharpoonright n}$ . Then  $\gamma$  is in  $M_\alpha \cap \omega_1 = \delta$ .

We prove that  $M$  is not in  $V$  by showing how to compute  $f$  by induction from  $M$ . Suppose  $f \upharpoonright n$  is known. Fix  $j < 2$  such that  $f(n) \neq j$ . We claim that  $\xi_{(f \upharpoonright n) \frown j}$  is not in  $M$ . Otherwise there is  $k > n$  such that  $\xi_{(f \upharpoonright n) \frown j}$  is in  $M_k$ . Fix  $\alpha$  in  $Z_{f \upharpoonright k}$ . Then  $\xi_{(f \upharpoonright n) \frown j}$  is in  $M_\alpha$ . But  $\alpha$  is in  $Z_{f \upharpoonright (n+1)}$ , contradicting Lemma 2.6(4). So  $f(n)$  is the unique  $i < 2$  such that  $\xi_{(f \upharpoonright n) \frown i}$  is in  $M$ . This completes the definition of  $f$  from  $M$ . Since  $f$  is not in  $V$ , neither is  $M$ .

Let  $\langle D_n : n < \omega \rangle$  enumerate the dense subsets of  $\mathbb{P}(S)$  lying in  $N$ . Inductively define a decreasing sequence  $\langle q_n : n < \omega \rangle$  in  $N \cap \mathbb{P}(S)$  such that  $q_0 = q$  and  $q_{n+1}$  is in  $D_n \cap N$ . Write  $\bigcup \{q_n : n < \omega\} = \langle b_i : i < \gamma \rangle$ . Clearly  $\bigcup \{b_i : i < \gamma\} = N \cap \omega_2 = M$ , and since  $M$  is not in  $V$ ,  $r = \langle b_i : i < \gamma \rangle \cup \{\langle \gamma, M \rangle\}$  is a condition in  $\mathbb{P}(S)$ . By an easy density argument,  $r$  forces that  $N \cap \omega_1 = \delta$  is a limit point of  $\dot{C}$ , and hence is in  $\dot{C}$ . Let  $\dot{r}$  be a name for  $r$ . Then  $p * \dot{r} \leq p * \dot{q}$  and  $p * \dot{r}$  forces  $\delta$  is in  $A \cap \dot{C}$ .  $\square$

The proof of Theorem 4.4 above is similar to the proof of Theorem 4.2.

Now we are ready to prove that MM implies there exists a disjoint club sequence on  $\omega_2$ .

*Proof of Theorem 4.1.* Assume Martin's Maximum. Inductively define  $A$  and  $\langle C_\alpha : \alpha \in A \rangle$  as follows. Suppose  $\alpha$  is in  $\omega_2 \cap \text{cof}(\omega_1)$  and  $A \cap \alpha$  and  $\langle C_\beta : \beta \in A \cap \alpha \rangle$  are defined. Let  $\alpha$  be in  $A$  iff the set  $\bigcup \{C_\beta : \beta \in A \cap \alpha\}$  is non-stationary in  $P_{\omega_1}(\alpha)$ .

If  $\alpha$  is in  $A$  then choose a club set  $\mathcal{C}_\alpha \subseteq P_{\omega_1}(\alpha)$  with size  $\omega_1$  which is disjoint from this union.

This completes the definition. We prove that  $A$  is stationary. Then clearly  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  is a disjoint club sequence. Fix a club set  $C \subseteq \omega_2$ .

Let  $\text{ADD}$  denote the forcing poset for adding a single Cohen real with finite conditions and let  $\dot{S}$  be an  $\text{ADD}$ -name for the set  $(P_{\omega_1}(\omega_2) \setminus V)$ . By Theorem 4.4 the poset  $\text{ADD} * \mathbb{P}(\dot{S})$  preserves stationary subsets of  $\omega_1$ . We will apply Martin's Maximum to this poset after choosing a suitable collection of dense sets.

For each  $\alpha < \omega_2$  fix a surjection  $f_\alpha : \omega_1 \rightarrow \alpha$ . If  $\beta$  is in  $A$  enumerate  $\mathcal{C}_\beta$  as  $\langle a_i^\beta : i < \omega_1 \rangle$ . For every quadruple  $i, j, k, l$  of countable ordinals let  $D(i, j, k, l)$  denote the set of conditions  $p * \dot{q}$  such that:

- (1)  $p$  forces that  $i$  and  $j$  are in the domain of  $\dot{q}$ , and for some  $\beta_i$  and  $\beta_j$ ,  $p$  forces  $\beta_i = \sup(\dot{q}(i))$  and  $\beta_j = \sup(\dot{q}(j))$ ,
- (2) there is some  $\zeta < \omega_1$  such that  $p$  forces  $\zeta$  is the least element in  $\text{dom}(\dot{q})$  such that  $f_{\beta_i}(j) \in \dot{q}(\zeta)$ ,
- (3) there is  $\xi$  in  $C$  larger than  $\beta_i$  and  $\beta_j$  such that  $p$  forces  $\xi$  is the supremum of the maximal set in  $\dot{q}$ ,
- (4) if  $f_{\beta_j}(k) = \gamma$  is in  $A$ , then there is  $z$  such that  $p$  forces  $z$  is in the symmetric difference  $\dot{q}(i) \Delta a_l^\gamma$ .

It is routine to check that  $D(i, j, k, l)$  is dense.

Let  $G * H$  be a filter on  $\text{ADD} * \mathbb{P}(\dot{S})$  intersecting each  $D(i, j, k, l)$ . For  $i < \omega_1$  define  $a_i$  as the set of  $\beta$  for which there exists some  $p * \dot{q}$  in  $G * H$  such that  $p$  forces  $i \in \text{dom}(\dot{q})$  and  $p$  forces  $\beta$  is in  $\dot{q}(i)$ . The definition of the dense sets implies that  $\langle a_i : i < \omega_1 \rangle$  is increasing, continuous, and cofinal in  $P_{\omega_1}(\alpha)$  for some  $\alpha$  in  $C \cap \text{cof}(\omega_1)$ . By (4), for each  $\gamma$  in  $A \cap \alpha$ ,  $\{a_i : i < \omega_1\}$  is disjoint from  $\mathcal{C}_\gamma$ . Therefore  $\bigcup \{\mathcal{C}_\gamma : \gamma \in A \cap \alpha\}$  is non-stationary in  $P_{\omega_1}(\alpha)$ , hence by the definition of  $A$ ,  $\alpha$  is in  $A \cap C$ . So  $A$  is stationary.  $\square$

## 5. THE EQUICONSISTENCY RESULT

We now prove Theorem 0.1 establishing the consistency strength of each of the following statements to be exactly a Mahlo cardinal: (1) There does not exist a thin stationary subset of  $P_{\omega_1}(\omega_2)$ . (2) There exists a disjoint club sequence on  $\omega_2$ . (3) There exists a fat stationary set  $S \subseteq \omega_2$  such that any forcing poset which preserves  $\omega_1$  and  $\omega_2$  does not add a club subset to  $S$ .

By [5] if there exists a thin stationary subset of  $P_{\omega_1}(\omega_2)$  then for any fat stationary set  $S \subseteq \omega_2$ , there is a forcing poset which preserves cardinals and adds a club subset to  $S$ . So (2) and (3) both imply (1), which in turn implies there is no special Aronszajn tree on  $\omega_2$ . So  $\omega_2$  is Mahlo in  $L$  by [8].

In the other direction assume that  $\kappa$  is a Mahlo cardinal. We will define a revised countable support iteration which collapses  $\kappa$  to become  $\omega_2$  and adds a disjoint club sequence on  $\omega_2$ . At individual stages of the iteration we force with either a collapse forcing or the poset  $\text{ADD} * \mathbb{P}(\dot{S})$  from the previous section. To ensure that  $\omega_1$  is not collapsed we verify that  $\text{ADD} * \mathbb{P}(\dot{S})$  satisfies an iterable condition known as the  $\mathbb{I}$ -universal property. Our description of this construction is self-contained, except for the proof of Theorem 5.9 which summarizes the relevant properties of the RCS iteration. For more information on such iterations and the  $\mathbb{I}$ -universal property see [10].

**Definition 5.1.** A pair  $\langle T, \mathbf{I} \rangle$  is a tagged tree if:

- (1)  $T \subseteq {}^{<\omega}\mathbf{On}$  is a tree such that each  $\eta$  in  $T$  has at least one successor,
- (2)  $\mathbf{I} : T \rightarrow V$  is a partial function such that each  $\mathbf{I}(\eta)$  is an ideal on some set  $X_\eta$  and for each  $\eta$  in the domain of  $\mathbf{I}$ , the set  $\{\alpha : \eta \hat{=} \alpha \in T\}$  is in  $(\mathbf{I}(\eta))^+$ ,
- (3) for each cofinal branch  $b$  of  $T$ , there are infinitely many  $n < \omega$  such that  $b \upharpoonright n$  is in the domain of  $\mathbf{I}$ .

If  $\eta$  is in the domain of  $\mathbf{I}$ , we say that  $\eta$  is a *splitting point* of  $T$ . It follows from (1) and (3) that for every  $\eta$  in  $T$  there is  $\eta \triangleleft \nu$  which is a splitting point.

**Definition 5.2.** Let  $\mathbb{I}$  be a family of ideals and  $\langle T, \mathbf{I} \rangle$  a tagged tree. Then  $\langle T, \mathbf{I} \rangle$  is an  $\mathbb{I}$ -tree if for each splitting point  $\eta$  in  $T$ ,  $\mathbf{I}(\eta)$  is in  $\mathbb{I}$ .

Suppose  $T \subseteq {}^{<\omega}\mathbf{On}$  is a tree. If  $\eta$  is in  $T$ , let  $T^{[\eta]}$  denote the tree  $\{\nu \in T : \nu \trianglelefteq \eta \text{ or } \eta \trianglelefteq \nu\}$ . A set  $J \subseteq T$  is called a *front* if for distinct nodes in  $J$ , neither is an initial segment of the other, and for any cofinal branch  $b$  of  $T$  there is  $\eta$  in  $J$  which is an initial segment of  $b$ .

**Definition 5.3.** Suppose  $\langle T, \mathbf{I} \rangle$  is tagged tree. Let  $\theta$  be a regular cardinal such that  $\langle T, \mathbf{I} \rangle$  is in  $H(\theta)$ , and let  $<_\theta$  be a well-ordering of  $H(\theta)$ . A sequence  $\langle N_\eta : \eta \in T \rangle$  is a tree of models for  $\theta$  provided that:

- (1) each  $N_\eta$  is a countable elementary substructure of  $\langle H(\theta), \in, <_\theta, \langle T, \mathbf{I} \rangle \rangle$ ,
- (2) if  $\eta \triangleleft \nu$  in  $T$ , then  $N_\eta \prec N_\nu$ ,
- (3) for each  $\eta$  in  $T$ ,  $\eta$  is in  $N_\eta$ .

**Definition 5.4.** Suppose  $\langle T, \mathbf{I} \rangle$  is an  $\mathbb{I}$ -tree, and  $\theta$  is a regular cardinal such that  $H(\theta)$  contains  $\langle T, \mathbf{I} \rangle$  and  $\mathbb{I}$ . A sequence  $\langle N_\eta : \eta \in T \rangle$  is an  $\mathbb{I}$ -suitable tree of models for  $\theta$  if it is a tree of models for  $\theta$  and for every  $\eta$  in  $T$  and  $I$  in  $\mathbb{I} \cap N_\eta$ , the set

$$\{\nu \in T^{[\eta]} : \nu \text{ is a splitting point and } \mathbf{I}(\nu) = I\}$$

contains a front in  $T^{[\eta]}$ .

**Definition 5.5.** Let  $\langle T, \mathbf{I} \rangle$ ,  $\mathbb{I}$ , and  $\theta$  be as in Definition 5.4. A sequence  $\langle N_\eta : \eta \in T \rangle$  is an  $\omega_1$ -strictly  $\mathbb{I}$ -suitable tree of models for  $\theta$  if it is an  $\mathbb{I}$ -suitable tree of models for  $\theta$  and there exists  $\delta < \omega_1$  such that for all  $\eta$  in  $T$ ,  $N_\eta \cap \omega_1 = \delta$ .

If  $\langle N_\eta : \eta \in T \rangle$  is a tree of models and  $b$  is a cofinal branch of  $T$ , write  $N_b$  for the set  $\bigcup \{N_{b \upharpoonright n} : n < \omega\}$ . Note that if  $\langle N_\eta : \eta \in T \rangle$  is an  $\omega_1$ -strictly  $\mathbb{I}$ -suitable tree of models for  $\theta$ , then for any cofinal branch  $b$  of  $T$ ,  $N_b \cap \omega_1 = N_\emptyset \cap \omega_1$ .

**Lemma 5.6.** Let  $\langle T, \mathbf{I} \rangle$ ,  $\mathbb{I}$ , and  $\theta$  be as in Definition 5.4, and let  $\langle N_\eta : \eta \in T \rangle$  be an  $\omega_1$ -strictly  $\mathbb{I}$ -suitable tree of models for  $\theta$ . Suppose  $\eta \triangleleft \nu$  in  $T$  and  $(N_\nu \cap \omega_2) \setminus N_\eta$  is non-empty. Let  $\gamma$  be the minimum element of  $(N_\nu \cap \omega_2) \setminus N_\eta$ . Then  $\gamma \geq \sup(N_\eta \cap \omega_2)$ .

*Proof.* Otherwise there is  $\beta$  in  $N_\eta \cap \omega_2$  such that  $\gamma < \beta$ . By elementarity, there is a surjection  $f : \omega_1 \rightarrow \beta$  in  $N_\eta$ . So  $f^{-1}(\gamma) \in N_\nu \cap \omega_1 = N_\eta \cap \omega_1$ . Hence  $f(f^{-1}(\gamma)) = \gamma$  is in  $N_\eta$ , which is a contradiction.  $\square$

Let  $\mathbb{I}$  be a family of ideals. We say that  $\mathbb{I}$  is *restriction-closed* if for all  $I$  in  $\mathbb{I}$ , for any set  $A$  in  $I^+$ , the ideal  $I \upharpoonright A$  is in  $\mathbb{I}$ . If  $\mu$  is a regular uncountable cardinal, we say that  $\mathbb{I}$  is  $\mu$ -complete if each ideal in  $\mathbb{I}$  is  $\mu$ -complete.

**Definition 5.7.** Suppose that  $\mathbb{I}$  is a non-empty restriction-closed  $\omega_2$ -complete family of ideals and let  $\mathbb{P}$  be a forcing poset. Then  $\mathbb{P}$  satisfies the  $\mathbb{I}$ -universal property

if for all sufficiently large regular cardinals  $\theta$  with  $\mathbb{I}$  in  $H(\theta)$ , if  $\langle N_\eta : \eta \in T \rangle$  is an  $\omega_1$ -strictly  $\mathbb{I}$ -suitable tree of models for  $\theta$ , then for all  $p$  in  $N_\langle \rangle \cap \mathbb{P}$  there is  $q \leq p$  such that  $q$  forces there is a cofinal branch  $b$  of  $T$  such that  $N_b[\dot{G}] \cap \omega_1 = N_\langle \rangle \cap \omega_1$ .

Definition 5.7 is Shelah's characterization of the  $\mathbb{I}$ -universal property given in [10] Chapter XV 2.11, 2.12, and 2.13. Note that in the definition,  $q$  is semigeneric for  $N_\langle \rangle$ . In 2.12 Shelah proves that there are stationarily many structures  $N$  for which  $N = N_\langle \rangle$  for some  $\omega_1$ -strictly  $\mathbb{I}$ -suitable tree of models  $\langle N_\eta : \eta \in T \rangle$ . So by standard arguments if  $\mathbb{P}$  satisfies the  $\mathbb{I}$ -universal property then  $\mathbb{P}$  preserves  $\omega_1$  and preserves stationary subsets of  $\omega_1$ . Note that any semiproper forcing poset satisfies the  $\mathbb{I}$ -universal property.

**Theorem 5.8.** *Let  $\mathbb{I}$  be the family of ideals of the form  $NS_{\omega_2} \upharpoonright A$ , where  $A$  is a stationary subset of  $\omega_2 \cap \text{cof}(\omega)$ . Let  $\dot{S}$  be an ADD-name for the set  $(P_{\omega_1}(\omega_2) \setminus V)$ . Then  $\text{ADD} * \mathbb{P}(\dot{S})$  satisfies the  $\mathbb{I}$ -universal property.*

*Proof.* Fix a regular cardinal  $\theta \gg \omega_2$  and let  $\langle N_\eta : \eta \in T \rangle$  be an  $\omega_1$ -strictly  $\mathbb{I}$ -suitable tree of models for  $\theta$ . Let  $p * \dot{q}$  be a condition in  $(\text{ADD} * \mathbb{P}(\dot{S})) \cap N_\langle \rangle$ . We find a refinement of  $p * \dot{q}$  which forces there is a cofinal branch  $b$  of  $T$  such that  $N_b[\dot{G} * \dot{H}] \cap \omega_1 = N_\langle \rangle \cap \omega_1$ .

We use an argument similar to the proof of Lemma 2.6 to define a sequence  $\langle \eta_s, \xi_s : s \in {}^{<\omega}2 \rangle$  satisfying:

- (1) each  $\eta_s$  is in  $T$ , each  $\xi_s$  is in  $N_{\eta_s} \cap \omega_2$ , and  $s \triangleleft t$  implies  $\eta_s \triangleleft \eta_t$ ,
- (2) if  $s \hat{\sim} 0 \trianglelefteq t$  then  $\xi_{s \hat{\sim} 1}$  is not in  $N_{\eta_t}$ , and if  $s \hat{\sim} 1 \trianglelefteq u$  then  $\xi_{s \hat{\sim} 0}$  is not in  $N_{\eta_u}$ .

Let  $\eta_\langle \rangle = \langle \rangle$  and  $\xi_\langle \rangle = 0$ . Suppose  $\eta_s$  is defined. Choose a splitting point  $\nu_s$  in  $T$  above  $\eta_s$ . Let  $Z$  denote the set of  $\alpha < \omega_2$  such that  $\nu_s \hat{\sim} \alpha$  is in  $T$ . Since  $\nu_s$  is a splitting point, by the definition of  $\mathbb{I}$  the set  $Z$  is a stationary subset of  $\omega_2 \cap \text{cof}(\omega)$ . For each  $\alpha$  in  $Z$ ,  $\alpha$  is in  $N_{(\nu_s \hat{\sim} \alpha)}$  and has cofinality  $\omega$ , so  $N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha$  is a countable cofinal subset of  $\alpha$ . Define  $X_s$  as the set of  $\xi$  in  $\omega_2$  such that the set

$$\{\alpha \in Z : \xi \in N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha\}$$

is stationary. An easy argument using Fodor's Lemma shows that  $X_s$  is unbounded in  $\omega_2$ . For all large enough  $\alpha$  in  $Z$ , the set  $(X_s \setminus \sup(N_{\nu_s} \cap \omega_2)) \cap \alpha$  has size  $\omega_1$ . So there is a stationary set  $Z'_1 \subseteq Z$  and an ordinal  $\xi_{s \hat{\sim} 0}$  in  $X_s$  such that  $\xi_{s \hat{\sim} 0}$  is larger than  $\sup(N_{\nu_s} \cap \omega_2)$  and for all  $\alpha$  in  $Z'_1$ ,  $\xi_{s \hat{\sim} 0}$  is not in  $N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha$ . Let  $Z'_0$  be the stationary set of  $\alpha$  in  $Z$  such that  $\xi_{s \hat{\sim} 0}$  is in  $N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha$ . Now define  $Y_s$  as the set of  $\xi$  in  $\omega_2$  such that the set

$$\{\alpha \in Z'_1 : \xi \in N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha\}$$

is stationary. Again we can find  $Z_0 \subseteq Z'_0$  stationary and  $\xi_{s \hat{\sim} 1}$  in  $Y_s$  such that  $\xi_{s \hat{\sim} 1}$  is larger than  $\sup(N_{\nu_s} \cap \omega_2)$  and for all  $\alpha$  in  $Z_0$ ,  $\xi_{s \hat{\sim} 1}$  is not in  $N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha$ . Let  $Z_1$  be the stationary set of  $\alpha$  in  $Z'_1$  such that  $\xi_{s \hat{\sim} 1}$  is in  $N_{(\nu_s \hat{\sim} \alpha)} \cap \alpha$ .

Now define  $\eta_{s \hat{\sim} 0}$  to be equal to  $\nu_s \hat{\sim} \alpha$  for some  $\alpha$  in  $Z_0$  larger than  $\xi_{s \hat{\sim} 1}$ , and define  $\eta_{s \hat{\sim} 1}$  to be  $\nu_s \hat{\sim} \beta$  for some  $\beta$  in  $Z_1$  larger than  $\xi_{s \hat{\sim} 0}$ . By definition  $\xi_{s \hat{\sim} 0}$  is in  $N_{\eta_{s \hat{\sim} 0}}$  and  $\xi_{s \hat{\sim} 1}$  is in  $N_{\eta_{s \hat{\sim} 1}}$ .

We claim that if  $\eta_{s \hat{\sim} 0} \trianglelefteq \nu$  in  $T$ , then  $\xi_{s \hat{\sim} 1}$  is not in  $N_\nu$ . Since  $\alpha$  is in  $Z_0$ ,  $\xi_{s \hat{\sim} 1}$  is not in  $N_{(\eta_{s \hat{\sim} 0})} \cap \alpha$ . But  $\xi_{s \hat{\sim} 1} < \alpha$ , so  $\xi_{s \hat{\sim} 1}$  is not in  $N_{(\eta_{s \hat{\sim} 0})}$ . By Lemma 5.6 the minimum element of  $N_\nu \cap \omega_2$  which is not in  $N_{(\eta_{s \hat{\sim} 0})}$ , if such an ordinal exists, is at least  $\sup(N_{(\eta_{s \hat{\sim} 0})} \cap \omega_2) \geq \alpha > \xi_{s \hat{\sim} 1}$ . So  $\xi_{s \hat{\sim} 1}$  is not in  $N_\nu$ . Similarly if  $\eta_{s \hat{\sim} 1} \trianglelefteq \nu$  in  $T$ , then  $\xi_{s \hat{\sim} 0}$  is not in  $N_\nu$ . This completes the definition. Conditions (1) and (2) are now easily verified.

Since  $\mathbb{P}$  is  $\omega_1$ -c.c., the condition  $p$  itself is generic for each  $N_\eta$ . Let  $G$  be a generic filter for ADD over  $V$  which contains  $p$ . Then for all  $\eta$  in  $T$ ,  $N_\eta[G] \cap \omega_2 = N_\eta \cap \omega_2$ . So for any cofinal branch  $b$  of  $T$  in  $V[G]$ ,  $N_b[G] \cap \omega_2 = \bigcup \{N_{b \upharpoonright n} \cap \omega_2 : n < \omega\}$ ; in particular,  $N_b[G] \cap \omega_1 = N_\emptyset \cap \omega_1$ .

Let  $f : \omega \rightarrow 2$  be a function in  $V[G] \setminus V$ . Define  $b_f = \bigcup \{\eta_{f \upharpoonright n} : n < \omega\}$ . We prove that  $N_{b_f} \cap \omega_2$  is not in  $V$  by showing how to define  $f$  inductively from this set. Suppose  $f \upharpoonright n$  is known. Fix  $j < 2$  such that  $f(n) \neq j$ . We claim that  $\xi^* = \xi_{(f \upharpoonright n) \frown j}$  is not in  $N_{b_f} \cap \omega_2$ . Otherwise there is  $k > n$  such that  $\xi^*$  is in  $N_{\eta_{f \upharpoonright k}}$ . But  $f \upharpoonright (n+1) \sqsubseteq f \upharpoonright k$ . So by condition (2),  $\xi^*$  is not in  $N_{\eta_{f \upharpoonright k}}$ , which is a contradiction. So  $f(n)$  is the unique  $i < 2$  such that  $\xi_{(f \upharpoonright n) \frown i}$  is in  $N_{b_f} \cap \omega_2$ .

Let  $\langle D_n : n < \omega \rangle$  enumerate all the dense subsets of  $\mathbb{P}(S)$  lying in  $N_{b_f}[G]$ . Inductively define a sequence  $\langle q_n : n < \omega \rangle$  by letting  $q_0 = q$  and choosing  $q_{n+1}$  to be a refinement of  $q$  in  $D_n \cap N_{b_f}[G]$ . Let  $\langle b_i : i < \gamma \rangle = \bigcup \{q_n : n < \omega\}$ . Clearly  $\bigcup \{b_i : i < \gamma\} = N_{b_f} \cap \omega_2$ . Since  $N_{b_f} \cap \omega_2$  is not in  $V$ ,  $r = \langle b_i : i < \gamma \rangle \frown (N_{b_f} \cap \omega_2)$  is a condition in  $\mathbb{P}(S)$  below  $q$  and  $r$  is generic for  $N_{b_f}[G]$ . So  $r$  forces  $N_{b_f}[G][\dot{H}] \cap \omega_1 = N_{b_f}[G] \cap \omega_1 = N_\emptyset \cap \omega_1$ . Let  $\dot{r}$  be a name for  $r$ . Then  $p * \dot{r} \leq p * \dot{q}$  is as required.  $\square$

We state without proof the facts concerning RCS iterations which we shall use. These facts follow immediately from [10] Chapter XI 1.13 and Chapter XV 4.15.

**Theorem 5.9.** *Suppose  $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \alpha, j < \alpha \rangle$  is an RCS iteration. Then  $\mathbb{P}_\alpha$  preserves  $\omega_1$  if the iteration satisfies the following properties:*

- (1) *for each  $i < \alpha$  there is  $n < \omega$  such that  $\mathbb{P}_{i+n} \Vdash |\mathbb{P}_i| \leq \omega_1$ ,*
- (2) *for each  $i < \alpha$  there is an uncountable regular cardinal  $\kappa_i$  and a  $\mathbb{P}_i$ -name  $\dot{\mathbb{I}}_i$  such that  $\mathbb{P}_i$  is  $\kappa_i$ -c.c. and  $\mathbb{P}_i$  forces  $\dot{\mathbb{I}}_i$  is a non-empty restriction-closed  $\kappa_i$ -complete family of ideals such that  $\dot{\mathbb{Q}}_i$  satisfies the  $\dot{\mathbb{I}}_i$ -universal property.*

**Theorem 5.10.** *Let  $\alpha$  be a strongly inaccessible cardinal. Suppose that  $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \alpha, j < \alpha \rangle$  is a revised countable support iteration such that  $\mathbb{P}_\alpha$  preserves  $\omega_1$  and for all  $i < \alpha$ ,  $|\mathbb{P}_i| < \alpha$ . Then  $\mathbb{P}_\alpha$  is  $\alpha$ -c.c.*

Suppose  $\kappa$  is a Mahlo cardinal and let  $A$  be the stationary set of strongly inaccessible cardinals below  $\kappa$ . Define an RCS iteration  $\langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \kappa, j < \kappa \rangle$  by recursion as follows. Our recursion hypotheses will include that each  $\mathbb{P}_\alpha$  preserves  $\omega_1$ , and is  $\alpha$ -c.c. if  $\alpha$  is in  $A$ .

Suppose  $\mathbb{P}_\alpha$  is defined. If  $\alpha$  is not in  $A$  then let  $\mathbb{Q}_\alpha$  be a name for  $\text{COLL}(\omega_1, |\mathbb{P}_\alpha|)$ . Suppose  $\alpha$  is in  $A$ . By the recursion hypotheses  $\mathbb{P}_\alpha$  forces  $\alpha = \omega_2$ . Let  $\dot{\mathbb{Q}}_\alpha$  be a name for the poset  $\text{ADD} * \mathbb{P}(\dot{S})$ .

If  $\alpha$  is not in  $A$  then choose some regular cardinal  $\kappa_\alpha$  larger than  $|\mathbb{P}_\alpha|$ , and let  $\dot{\mathbb{I}}_\alpha$  be a name for some non-empty restriction-closed  $\kappa_\alpha$ -complete family of ideals on  $\kappa_\alpha$ . Then  $\mathbb{P}_\alpha$  is  $\kappa_\alpha$ -c.c., and since  $\dot{\mathbb{Q}}_\alpha$  is proper,  $\mathbb{P}_\alpha$  forces  $\dot{\mathbb{Q}}_\alpha$  satisfies the  $\dot{\mathbb{I}}_\alpha$ -universal property. Suppose  $\alpha$  is in  $A$ . Then let  $\alpha = \kappa_\alpha$  and define  $\dot{\mathbb{I}}_\alpha$  as a name for the family of ideals on  $\omega_2$  as described in Theorem 5.8. Then  $\mathbb{P}_\alpha$  is  $\kappa_\alpha$ -c.c. and forces  $\dot{\mathbb{Q}}_\alpha$  satisfies the  $\dot{\mathbb{I}}_\alpha$ -universal property.

Suppose  $\beta \leq \kappa$  is a limit ordinal and  $\mathbb{P}_\alpha$  is defined for all  $\alpha < \beta$ . Define  $\mathbb{P}_\beta$  as the revised countable support limit of  $\langle \mathbb{P}_\alpha : \alpha < \beta \rangle$ . By Theorem 5.9 and the recursion hypotheses,  $\mathbb{P}_\beta$  preserves  $\omega_1$ . Hence if  $\beta$  is in  $A \cup \{\kappa\}$ , then  $\mathbb{P}_\beta$  is  $\beta$ -c.c. by Theorem 5.10.

This completes the definition. Let  $G$  be generic for  $\mathbb{P}_\kappa$ . The poset  $\mathbb{P}_\kappa$  is  $\kappa$ -c.c. and preserves  $\omega_1$ , so in  $V[G]$  we have that  $\kappa = \omega_2$  and  $A$  is a stationary subset of

$\omega_2 \cap \text{cof}(\omega_1)$ . For each  $\alpha$  in  $A$  let  $\mathcal{C}_\alpha$  be the club on  $P_{\omega_1}(\alpha)$  introduced by  $\mathbb{Q}_\alpha$ . If  $\alpha < \beta$  are in  $A$ , then  $\mathcal{C}_\alpha$  and  $\mathcal{C}_\beta$  are disjoint since  $\mathcal{C}_\beta$  is disjoint from  $V[G \upharpoonright \beta]$ . So  $\langle \mathcal{C}_\alpha : \alpha \in A \rangle$  is a disjoint club sequence on  $\omega_2$  in  $V[G]$ .

We conclude the paper with several questions.

(1) Assuming Martin's Maximum, the poset  $\text{ADD} * \mathbb{P}(\dot{S})$  is semiproper. Is this poset semiproper in general?

(2) Is it consistent that there exists a stationary set  $A \subseteq \omega_2 \cap \text{cof}(\omega_1)$  such that neither  $A \cup \text{cof}(\omega)$  nor  $\omega_2 \setminus A$  can acquire a club subset in an  $\omega_1$  and  $\omega_2$  preserving extension?

(3) To what extent can the results of this paper be extended to cardinals greater than  $\omega_2$ ? For example, is it consistent that there is a fat stationary subset of  $\omega_3$  which cannot acquire a club subset by any forcing poset which preserves  $\omega_1$ ,  $\omega_2$ , and  $\omega_3$ ?

#### REFERENCES

- [1] U. Abraham and S. Shelah. Forcing closed unbounded sets. *Journal of Symbolic Logic*, 48(3):643–657, 1983.
- [2] J. Baumgartner and A. Taylor. Saturation properties of ideals in generic extensions I. *Transactions of the American Mathematical Society*, 270(2):557–574, 1982.
- [3] Q. Feng, T. Jech, and J. Zapletal. On the structure of stationary sets. Preprint.
- [4] M. Foreman, M. Magidor, and S. Shelah. Martin's maximum, saturated ideals, and non-regular ultrafilters. part I. *Annals of Mathematics*, 127:1–47, 1988.
- [5] S. Friedman. Forcing with finite conditions. Preprint.
- [6] M. Gitik. Nonsplitting subset of  $P_\kappa \kappa^+$ . *J. Symbolic Logic*, 50(4):881–894, 1985.
- [7] W. Mitchell.  $I[\omega_2]$  can be the non-stationary ideal. Preprint.
- [8] W. Mitchell. Aronszajn trees and the independence of the transfer property. *Annals of Mathematical Logic*, pages 21–46, 1972.
- [9] M. Rubin and S. Shelah. Combinatorial problems on trees: Partitions,  $\Delta$ -systems and large free subtrees. *Annals of Pure and Applied Logic*, 33:43–81, 1987.
- [10] S. Shelah. *Proper and Improper Forcing*. Springer, 1998.

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, UNIVERSITY OF VIENNA  
*E-mail address:* `sdf@logic.univie.ac.at`  
*URL:* `http://www.logic.univie.ac.at/~sdf`

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, UNIVERSITY OF VIENNA  
*E-mail address:* `jkruieger@logic.univie.ac.at`  
*URL:* `http://www.logic.univie.ac.at/~jkruieger`